

EPA Jacket 228-528

Vol.4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 27, 2015

Carrie M. Tackema
Regulatory Manager
Nufarm Americas, Inc.
11901 S. Austin Avenue
Alsip, IL 60803

Subject: CSF Amendment –New Alternate CSF 9
Product Name: Nuprid 4F Insecticide
EPA Registration Number: 228-528
Application Date: 10/06/14
Decision Number: 497329

Dear Carrie M. Tackema:

The Confidential Statement of Formula (CSF) referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

Please note that the record for this product currently contains the following CSFs:

- Basic CSF dated 07/02/2012
- Alternate CSF 1 dated 10/10/2012
- Alternate CSF 2 dated 10/10/2012
- Alternate CSF 3 dated 10/10/2012
- Alternate CSF 4 dated 10/10/2012
- Alternate CSF 5 dated 10/10/2012
- Alternate CSF 6 dated 10/10/2012
- Alternate CSF 7 dated 10/10/2012
- Alternate CSF 8 dated 10/10/2012
- Alternate CSF 9 dated 10/01/2014

Any CSFs other than those listed above are superseded/no longer valid. If you have any questions, please contact Jessica Rogala by phone at 703-347-0263, or via email at rogala.jessica@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'Mark Suarez', with a long horizontal stroke extending to the right.

Mark Suarez, Product Manager 07
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEE

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION DIVISION (7505P)

DP BARCODE No.: D 424142, REG. No.: 228-528, PRODUCT NAME: Nuprid 4 F
Insecticide Code(s): 129099, Decision: 497131, Action Code: R340, Food Use: [Yes]

DATE OUT: 01/26/15

SUBJECT: End Use Product Chemistry Review
Product Name: Nuprid 4F Insecticide

FROM: Indira Gairola, Product Chemistry Team
CITAB/ Registration Division (7505P)

(d) 01/26/15

SBur 01/26/15

TO: Gene Benbow / Venus Eagle PM 01
Insecticide Branch / Registration Division (7505P)

Company Name: Nufarm Americas, Inc.
Formulation Type: Herbicide

INTRODUCTION:

The registrant is submitting Alternate Formulation CSF #10 dated 11/04/14 for the subject product "Nuprid 4F Insecticide" along with Alternate formulation for the end use product "Nuprid 4F Insecticide".

CITAB (Chemistry Team) has been asked to determine the acceptability of the aforementioned CSF.

SUMMARY OF FINDINGS:

1. The alternate CSF dated 11/04/14 was reviewed for the subject product "Nuprid 4F Insecticide".
2. The certified limits are in compliance with standard certified limits.
3. Inerts used are approved for the proposed food use.
4. It is determined that the alternate CSF dated 11/04/14 is acceptable.

CONCLUSIONS:

CITAB/ Product Chemistry Team has reviewed the alternate CSF #10 dated 11/04/14 for the end use product "Nuprid 4F Insecticide" and has concluded:

1. The aforementioned alternate CSF for the subject product "Nuprid 4F Insecticide" is determined to be acceptable and will be placed in file for record.

21-Day Screen of Amendment
(Completed by Contractor)

21-day Expires on 11-27-14

Document Part Of: 228-528
MRID, If Any: 495030

Content Screen: Recommended to
Pass/Fail

11-3 Review: Passed/Failed/NA

Overall Status: Pass/Fail

Document returned to:

STEPHEN SUTABLE

E-SUBMISSION**Memorandum**Date: 11 / 14 / 14To: PMI, Regulatory Manager

From: Information Services Branch, ITRMD

Your receipt of this data submission is not an indication that MRIDs for the enclosed studies have been posted to OPPIN.

We expect that it will be approximately 5 days from the above date before the study-level data is available in OPPIN.

If you have any questions about this process, please contact Teresa Downs (305-5363).

This is a: ☒ fully accepted submission
☐ partially accepted submission
☐ rejected submission

PRIA 3 – 21 Day Content Screen Review Worksheet

(EPA/OPP Use Only)

September 2012

21 Day Screen Start Date: 11-6-14

Experts In-Processing Signature: B.O.

Date 11-10-14

Fee Paid: Yes ☒

Division management contacted on issues No ☐ Yes ☐ Date _____

EPA Reg. Number: <u>228-528</u>		EPA Receipt Date: <u>11-6-14</u>				
Items for Review				Yes	No	N/A*
1	Application Form (EPA Form 8570-1) signed & complete including package type			X		
2	Confidential Statement of Formula all boxes completed, form signed, and dated (EPA Form 8570-4)			X		
	a) All <u>inerts</u> , including fragrances, approved for the proposed uses (see Footnote A)	yes	no			
		X				
3	Certification with Respect to Citation of Data (EPA Form 8570-34) completed and signed (N/A if 100% repack)			X		
	Certificate and data matrix consistent			X		
	If applicant is relying on data that are compensable, is the offer to pay statement included. (see Footnote B)	yes	no			
	If applicable, is there a letter of Authorization for exclusive use only.					
4	Formulator's Exemption Statement (EPA Form 8570-27) completed and signed (N/A if source is unregistered or applicant owns the technical)					X
	Data Matrix (EPA Form 8570-35) both internal and external copies (PR 98-5) completed and signed (N/A if 100% repack)			X		
5	a) Selective Method (Fee category experts use)	yes	no			
		X				
	b) Cite-All (Fee category experts use)					
	c) Applicant owns all data (Fee category experts use)					
6	5 Copies of Label (Electronic labels on CD are encouraged and guidance is available)					X
7	Is the data package consistent with PR Notice 86-5			X		
8	Notice of Filing included with petitions					X

9	If applicable for conventional applications, <u>reduced risk rationale</u>			X
10	<u>Required Data</u> and/or data waivers. See Footnote C.			
	a) List study (or studies) not included with application			

Comments:

* Documentation: PASS / FAZL
 - required forms are complete

* Inerts: PASS / FAZL
 - approved for food use under 40 CFR 180.910, Pre & Post Harvest application

* PRN 11-3 PASS / FAZL
 - MRZD: 495030

* Overall Status: PASS / FAZL

JK 11-17-14

* N/A – Not Applicable

Footnotes

A. During the 21 day initial content review, all CSFs will be reviewed to determine whether all inerts listed, including fragrances, are approved for the proposed uses or have an application pending with the Agency. If an unapproved inert with no application pending with the Agency is identified, the applicant must either 1) resolve the inert issue by, for example, removing the inert, substituting it with an approved inert, submitting documentation that EPA approved the inert for the proposed pesticidal uses, correcting mistakes on the CSF, etc. or 2) provide the data to support OPP approval of the inert or 3) withdraw the application. Removing or substituting an inert ingredient will require a new CSF and may require submission of data. All information, forms, data and documentation resolving the inert issue must have been received by the Agency or the application withdrawn within the 21 day period, otherwise, the Agency will reject the application as described below.

To successfully complete this aspect of the 21 day initial content screen, applicants are **strongly encouraged** to verify that all inert ingredients have been approved for the application's uses or have an application pending with the Agency **even if a product is currently registered** by consulting the [inert Web site](#) and if the inert is not approved nor has an application pending with the Agency, to **obtain the necessary inert approval prior to submitting an application to register a pesticide product containing that inert ingredient**. Some inert ingredients are no longer approved for food uses or certain types of uses. The name and/or CAS number on a CSF must match the name and CAS number on this web site. Simple typographical errors in the name or CAS number have resulted in processing delays.

If an inert is not listed on the inert ingredient web site and the applicant believes that the inert has been approved, the applicant should contact the Inert Ingredient Assessment Branch (IIAB) at inertsbranch@epa.gov and resolve the issue. Copies of the correspondence with IIAB resolving the issue should accompany the application. All new inerts except PIP inerts are reviewed by IIAB. The IIAB should also be contacted for any questions on what supporting data needs to be submitted for and the Agency's inert review process. Questions on PIP inerts should be directed to the [Chief of Microbial Pesticides Branch](#).

When a brand, trade, or proprietary name of an inert ingredient is listed on a CSF, additional information such as an alternate name of the inert, CAS number or other information must also be included to enable the Agency to determine if it has been approved. Each component of an inert mixture (including a fragrance) must be identified. In some cases, the supplier of the mixture or fragrance may need to provide this information to the Agency. Prior to the Agency's receipt of an application, applicants must arrange with a proprietary mixture or fragrance supplier to provide the component information to the Agency or promptly upon EPA's request. If the inert ingredients in a proprietary blend (including fragrances) cannot or are not identified or provided within the 21-day content review period, the Agency will reject the application.

During the 21 day content review, applicants should submit information to the individual identified by the Agency when the applicant is informed of an unapproved inert.

Unapproved Inerts Identified on CSFs

All applications except conventional new products and PIPs

Once an unapproved inert is identified on a CSF, the Agency will contact the applicant with the following options:

1. Correct the application by, for instance, correcting the inert's identity or CAS number, providing documentation that the inert has been approved, or removing the unapproved inert from the CSF or replacing it with one that is approved for the application's uses; or
2. Provide the required information necessary to identify an inert approval application that is pending with the Agency; or
3. Submit the information and data needed for the Agency to approve the unapproved inert. If this option is selected and implemented, the Agency may request an extension in the PRIA decision review timeframe to accommodate the inert review/approval process;
4. Withdraw the application (the Agency retains 25% of the full fee for the fee category estimated); or

If none of these options is selected and implemented by the applicant within the 21 day content review period, the Agency will reject the application and retain 25% of the full fee of the category identified.

Conventional New Product Applications

When the Registration Division identifies an unapproved inert on a CSF with an application for a new product that the applicant has not identified as requiring an inert approval (R300 or R301), it will contact the applicant with the following options:

1. Correct the application by, for instance, correcting the inert's identity or CAS number, providing documentation that the inert has been approved, or removing the unapproved inert from the CSF or replacing it with one that is approved for the application's uses; or
2. Submit the information and data needed for the Agency to approve the unapproved inert, including any required petition to establish or amend a tolerance or exemption from a tolerance. (This option may change the PRIA category for the application, which could require a longer decision review time and a larger fee. If additional fees are due, they must be received by the Agency within the 21 day content review period.)

3. Withdraw the application (the Agency retains 25% of the full fee for the fee category estimated); or

If none of the above options is selected and implemented during the 21-day content-review period, the Agency will reject the application and retain 25% of the appropriate fee for the new product-inert approval category.

PIP Applications

When the Biopesticide and Pollution Prevention Division identifies an unapproved inert on a PIP CSF and a request to approve the inert does not accompany the application, it will contact the applicant with the following options:

1. Correct the application by, for instance, correcting the spelling or name of the inert to that in 40 CFR 174, or providing documentation that the inert has been approved; or
2. Submit the information and data needed for the Agency to approve the unapproved inert. If an inert ingredient tolerance exemption petition is required, the petition must be received by the Agency and the B903 fee paid within the 21 day period. If this option is selected and implemented, the Agency will discuss harmonizing the timeframe for both actions.
3. Withdraw the application (the Agency retains 25% of the full fee for the fee category estimated); or

If none of the above options is selected and implemented during the 21 day content review period, the Agency will reject the application and retain 25% of the fee.

B. A policy on documentation of offers to pay is still being developed, however, for a me-too or fast track (similar/identical) new product, R300 or A530, an application without the necessary authorizations of offers to pay will be placed into either R301 or A531. The Agency recommends that authorizations of offers to pay be submitted with other PRIA applications to avoid delays in the Agency's decision.

C. Biopesticide applicants are advised to contact the Agency and discuss study waivers prior to submitting their application to the Agency. Documentation of such discussions should be submitted with the study waiver.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

November 07, 2014

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

NUFARM AMERICAS, INC.
NUFARM AMERICAS, INC.
4020 AERIAL CENTER PKWY., STE. 101
MORRISVILLE, NC 27560

Report of Analysis for Compliance with PR Notice 11-03

Thank you for your submittal of 05-NOV-14. Our staff has completed a preliminary analysis of the material. The results are provided as follows:

Your submittal was found to be in full compliance with the standards for submission of data contained in PR Notice 11-03. A copy of your bibliography is enclosed, annotated with Master Record ID's (MRIDs) assigned to each document submitted. Please use these numbers in all future references to these documents. Thank you for your cooperation. If you have any questions concerning this data submission, please raise them with the cognizant Product Manager, to whom the data have been released.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

November 7, 2014

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

OPP Decision Number: D-497131
EPA File Symbol or Registration Number: 228-528
Product Name: NUPRID 4F INSECTICIDE
EPA Receipt Date: 05-Nov-2014
EPA Company Number: 228
Company Name: NUFARM AMERICAS, INC.

CARRIE M. TACKEMA
NUFARM AMERICAS, INC.
4020 AERIAL CENTER PKWY., STE. 101
MORRISVILLE, NC 27560-

SUBJECT: Receipt of Registration Amendment Subject to Registration Service Fee

Dear Registrant:

The Office of Pesticide Programs has received your amendment and certification of payment. If you submitted data with this application, the results of the PRN-2011-3 screen will be communicated separately. During the administrative screen, the Office of Pesticide Programs has determined that this Action is subject to a Pesticide Registration Service Fee as defined in the Pesticide Registration Improvement Act.

The Action has been identified as Action Code: R340

AMENDMENT;NON-FAST TRACK;REVIEW WITHIN RD, E.G. PRECAUTIONARY LABELING;

No additional payment is due at this time.

If you have any questions, please contact the Pesticide Registration Service Fee Ombudsman at (703) 308-9362.

Sincerely,

A handwritten signature in black ink, appearing to be "J. Sh".

Front End Processing Staff
Information Technology & Resources Management Division

Fee for Service

W
{960162A~

This package includes the following

- ☐ New Registration
- ☒ Amendment

☒ Studies? ☐ Fee Waiver?

☐ volpay % Reduction: ____

for Division

- ☐ AD
- ☐ BPPD
- ☒ RD

Risk Mgr. 1

Receipt No.

S-

960162

EPA File Symbol/Reg. No.

228-528

Pin-Punch Date:

11/6/2014

☐ This item is NOT subject to FFS action.

Action Code:

Requested:

R340

Granted:

R340

Amount Due: \$ 3,758.⁰⁰

Parent/Child Decisions:

☒ Inert Cleared for Intended Use

☐ Uncleared Inert in Product

Reviewer: 

Date: 11-7-14

Remarks:

e-Submission



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date November 4, 2014

EPA Reg. No./File Symbol: 228-528

Page 1 of 3

Applicant's/Registrant's Name & Address:

Nufarm Americas Inc.
11901 South Austin Ave.
Alsip, IL 60803

Product Name:

NUPRID 4F INSECTICIDE

Ingredient(s): Imidacloprid (PC Code 129099)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter (Company No.)	Status	Note
	PRODUCT CHEMISTRY DATA REQUIREMENTS				
830.1550	Product Identity and Composition	47042201	NUFARM (228)	OWN	
830.1600	Description of the Materials Used to Produce the Product	47042201	NUFARM (228)	OWN	
830.1620	Description of the Production Process	N/A			
830.1650	Description of the Formulation Process	47042201	NUFARM (228)	OWN	
830.1670	Discussion of the Formation of Impurities	47042201	NUFARM (228)	OWN	
830.1700	Preliminary Analysis	NA			
830.1750	Certified Limits	47042201	NUFARM (228)	OWN	
830.1800	Enforcement Analytical Method	47042201	NUFARM (228)	OWN	
830.6302	Color	47042202	NUFARM (228)	OWN	
830.6303	Physical State	47042202	NUFARM (228)	OWN	
830.6304	Odor	47042202	NUFARM (228)	OWN	
830.6313	Stability to normal / elevated temperatures, metals and metal ions	N/A			
830.6314	Oxidizing/Reducing Reaction	N/A			
830.6315	Flammability	N/A			
830.6316	Explosibility	N/A			

Signature

Name and Title:

Carrie M. Tackema
Regulatory Manager

Date

November 4, 2014



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date November 4, 2014

EPA Reg. No./File Symbol: 228-528

Page 2 of 3

Applicant's/Registrant's Name & Address:

Nufarm Americas Inc.
11901 South Austin Ave.
Alsip, IL 60803

Product Name:

NUPRID 4F INSECTICIDE

Ingredient(s): Imidacloprid (PC Code 129099)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter (Company No.)	Status	Note
830.6317	Storage Stability	47536501	NUFARM (228)	OWN	
830.6319	Miscibility	N/A			
830.6320	Corrosion Characteristics	47536501	NUFARM (228)	OWN	
830.6321	Dielectric Breakdown Voltage	N/A			
830.7000	pH	47042202	NUFARM (228)	OWN	
830.7050	UV/Visible Absorption	N/A			
830.7100	Viscosity	47042202	NUFARM (228)	OWN	
830.7200	Melting Point	N/A			
830.7220	Boiling Point	N/A			
830.7300	Density, Bulk Density, Specific Gravity	47042202	NUFARM (228)	OWN	
830.7370	Dissociation Constant	N/A			
830.7520	Particle Size	N/A			
830.7550 830.7560 830.7570	Partition Coefficient (n-octanol/water)	N/A			
830.7840 830.7860	Water Solubility	N/A			

Signature

Name and Title:

Carrie M. Tackema
Regulatory Manager

Date

November 4, 2014



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

401 M Street, S.W.
WASHINGTON, D.C. 20460

Form Approved OMB No. 2070-0060

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 0.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

DATA MATRIX

Date November 4, 2014

EPA Reg. No./File Symbol: 228-528

Page 3 of 3

Applicant's/Registrant's Name & Address:

Nufarm Americas Inc.
11901 South Austin Ave.
Alsip, IL 60803

Product Name:

NUPRID 4F INSECTICIDE

Ingredient(s): Imidacloprid (PC Code 129099)

Guideline Reference Number	Guideline Study Name	MRID Number	Submitter (Company No.)	Status	Note
830.7950	Vapor Pressure	N/A			
ACUTE TOXICITY DATA REQUIREMENTS					
870.1100 / 81-1	Acute Oral Toxicity (RAT)	47003904	NUFARM (228)	OWN	
870.1200 / 81-2	Acute Dermal Toxicity	47003905	NUFARM (228)	OWN	
870.1300 / 81-3	Acute Inhalation Toxicity	47003906	NUFARM (228)	OWN	
870.2400 / 81-4	Primary Eye Irritation	47003907	NUFARM (228)	OWN	
870.2500 / 81-5	Primary Skin Irritation	47003908	NUFARM (228)	OWN	
870.2600 / 81-6	Skin Sensitization	47003909	NUFARM (228)	OWN	

Generic Data Requirements: Cite-All - OTP

Syngenta Crop Protection, Inc. (100)	Bayer Healthcare LLC (011556)	United Phosphorus, Inc. (69811)	Rotam Agrochemical (83100)
Bayer CropScience LP (264)	Mitsui Chemicals, Inc. (33657)	Bayer Advanced (72155)	Sharda USA LLC (83529)
FMC Corp. Agricultural Products Group (279)	Loveland Products, Inc. (34704)	Scimetrics, Ltd. Corp. (72500)	Celsius Property BV (83558)
Chemtura Corporation (400)	Lanxess Corporation (39967)	Zelam Ltd. (72616)	Amtide (83851)
Bayer Environmental Sciences (432)	Makhteshim-Agan of North America, Inc. (66222)	Rockwell Laboratories Ltd. (73079)	Ensyslex IV (83923)
The Scotts Company (538)	Albaugh, Inc. (42750)	Sapstain Industry Group (73154)	Dalian Chemphy Chemicals (73467)
PBI Gordon (2217)	Aerixon, Inc. (43419)	Univar USA Inc. (73748)	
Bayer Corporation (3125)	Control Solutions (53883)	Innovative Pest Control Products (73766)	
Helena Chemical Company (5905)	Cheminova, Inc. (67760)	Arborjet (74578)	
Gustafson LLC (7501)	Mycogen Seeds (68467)	Arch Treatment Technologies, Inc. (75506)	
JJ Mauget Company (7946)	Arbor Systems, Inc. (69117)	Gro-Pro LLC (79676)	
The Andersons Lawn Fertilizer Division (9198)	Pet Logic, LLC (69332)	Rotam Ltd. (81598)	
Chemical Specialties, Inc. (10356)	Repar Corp. (69361)	Ensyslex III Inc. (82957)	

Signature

Name and Title:

Carrie M. Tackema
Regulatory Manager

Date

November 4, 2014



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
401 M Street, S.W.
WASHINGTON, D.C. 20460

Paperwork Reduction Act Notice: The public reporting burden for this collection of information is estimated to average 1.25 hours per response for registration activities and 0.25 hours per response for reregistration and special review activities, including time for reading the instructions and completing the necessary forms. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to: Director, OPPE Information Management Division (2137), U.S. Environmental Protection Agency, 401 M Street, S.W., Washington, DC 20460. Do not send the form to this address.

Certification with Respect to Citation of Data

Applicant's/Registrant's Name, Address and Telephone Number Nufarm Americas, Inc. 11901 South Austin Ave., Alsip, IL 60803	EPA Registration Number/ File Symbol 228-528
Active Ingredient(s) and/or representative test compound(s): Imidacloprid	Date 11-4-14
General use pattern(s) (list all those claimed for this product using 40 CFR Part 158) Terrestrial food/non-food crop; aquatic food/non-food crop; forestry; greenhouse food/non-food crop	Product Name Nuprid 4F Insecticide

NOTE: If your product is a 100% repackaging of another purchased EPA-registered product labeled for all the same uses on your label, you do not need to submit this form. You must submit the Formulator's Exemption Statement (EPA Form 8570-27).

- ☐ I am responding to a Data Call-In Notice, and have included with this form a list of companies sent offers of compensation (the Data Matrix form should be used for this purpose).

SECTION I: METHOD OF DATA SUPPORT (Check one method only)

- ☐ I am using the cite-all method of support, and have included with this form a list of companies sent offers of compensation (the Data Matrix form should be used for this purpose).
- ☒ I am using the selective method of support (or cite-all option under the selective method), and have included with this form a completed list of data requirements (the Data Matrix form must be used).

SECTION II: GENERAL OFFER TO PAY

[Required if using the cite-all method or when using the cite-all option under the selective method to satisfy one or more data requirements]

- ☒ I hereby offer and agree to pay compensation, to other persons, with regard to the approval of this application, to the extent required by FIFRA.

SECTION III: CERTIFICATION

I certify that this application for registration, this form for reregistration, or this Data Call-In Notice is supported by all data submitted or cited in the application for registration, the form for reregistration, or this Data Call-In response. In addition, if cite-all option under the selective method is indicated in Section I, this application is supported by all data in the Agency's files that (1) concern the properties or effects of this product or an identical or substantially similar product, or one or more of the ingredients in this product; and (2) is a type of data that would be required to be submitted under the data requirements in effect on the date of approval of this application if the application sought the initial registration of a product of identical or similar composition and uses.


I certify that for each exclusive use study cited in support of this registration or reregistration, that I am the original submitter or that I have obtained the written permission of the original submitter to cite that study.

I certify that for each study cited in support of this registration or reregistration that is not an exclusive use study, either: (a) I am the original data submitter; (b) I have obtained the written permission of the original data submitter to use this study in support of this application; (c) all periods of eligibility for compensation have expired for the study; (d) the study is in the public literature; or (e) I have notified in writing the company that submitted the study and have offered (i) to pay compensation to the extent required by sections 3(c)(1)(F) and/or 3(c)(2)(B) of FIFRA; and (ii) to commence negotiations to determine the amount and terms of compensation, if any, to be paid for the use of the study.

I certify that in all instances where an offer of compensation is required, copies of all offers to pay compensation and evidence of their delivery in accordance with sections 3(c)(1)(F) and/or 3(c)(2)(B) of FIFRA are available and will be submitted to the agency upon request. Should I fail to produce such evidence to the Agency upon request, I understand that the Agency may initiate action to deny, cancel or suspend the registration of my product in conformity with FIFRA.

I certify that the statements I have made on this form and all attachments to it are true, accurate and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Signature 	Date 11-4-14	Typed or Printed Name and Title Carrie M. Tackema, Regulatory Manager
---------------	------------------------	---



Receipt

Your payment is complete

Pay.gov Tracking ID: 25I9SGNG

Agency Tracking ID: 74702548829

Form Name: Pesticide Registration Improvement Act - Prepayment

Application Name: PRIA Service Fees

Payment Information

Payment Type: Debit or credit card

Payment Amount: \$3798.00

Transaction Date: 11/04/2014 11:51:47 AM EST

Payment Date: 11/04/2014

Registration Number: 228-528

Company Name: Nufarm Americas, Inc.

Company Number: 228

Action Code: R340

Account Information

Card Holder Name: Carrie Tackema

Billing Address: [REDACTED]

Billing Address 2: [REDACTED]

City: [REDACTED]

Country: United States

State/Province: [REDACTED]

ZIP/Postal Code: [REDACTED]

Card Type: American Express

Card Number: *****1008


Email Confirmation Receipt

Confirmation Receipts have been emailed to:

carrie.tackema@us.nufarm.com

Personal privacy information

e-Submission



8

Fee for Service

{958774V~

This package includes the following

- ☐ New Registration
- ☒ Amendment

☐ Studies? ☐ Fee Waiver?
☐ volpay % Reduction: ____

for Division

- ☐ AD
- ☐ BPPD
- ☒ RD

Risk Mgr. 1

Receipt No.

S- 958774

EPA File Symbol/Reg. No.

228-528

Pin-Punch Date:

10/7/2014

☒ This item is NOT subject to FFS action.

Action Code:

Requested:

Granted:

Amount Due: \$ _____

Parent/Child Decisions:

Rein 11-7-2014

☒ Inert Cleared for Intended Use

Uncleared Inert in Product

Reviewer: *J. GARVINE*

Date: *10/8/14*

Remarks:

S: 958774 Milestone Email:

Regulatory Type: Product Registration - Section 3

Resubmission: ☐ Yes ☒ No

Application Type: Amendment

Fee For Service: ☐ Yes ☒ No

Billable: ☐ Yes ☒ No

Company: 228 NUFARM AMERICAS, INC.



Print Letter

Enter More Information

Tracking

Risk Manager: Registration Division, Risk Management Team 1

Product #: 228-528 Product Name: NUPRID 4F INSECTICIDE

Override#:

Me Too Section3: 264-855

Me Too Product Name: TRIMAX PRO INSECTICIDE

Application Date: 01-Oct-2014



OPP Rec'd Date: 07-Oct-2014



Front End Date: 07-Oct-2014



Risk Manager Send Date: 08-Oct-2014



FFS Due Date:

Negotiated Due Date:

OPP Target Date:

Fast Track: ☐

New Ingredient: ☐

Receipt Description:

AMENDMENT

Receipt Content

CSF

View/Edit

Form A: ☐

Signature Date:

Form B: ☐

New Ingredient

Request Date:

New Ingredient

Received Date:

Signature Date:



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

October 8, 2014

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

MR. NATHAN P. EHRESMAN
NUFARM AMERICAS, INC.
NUFARM AMERICAS, INC.
4020 AERIAL CENTER PKWY., STE. 101
MORRISVILLE, NC 27560-

PRODUCT NAME: NUPRID 4F INSECTICIDE
COMPANY NAME: NUFARM AMERICAS, INC.
OPP IDENTIFICATION NUMBER:
EPA FILE SYMBOL: 228-528
EPA RECEIPT DATE: 10/07/14

SUBJECT: RECEIPT OF AMENDMENT

DEAR REGISTRANT:

The Office of Pesticide Programs has received your application for an amendment and it has passed an administrative screen for completeness.

During the initial screen we determined that the application appears to qualify for fast track review. The package will now be forwarded to the Product Manager for review to determine its acceptability for fast track status.

If you have any questions, please contact Registration Division, Risk Management Team 1, at (703) 308-8045.

Sincerely,

A handwritten signature in black ink, appearing to be "S. J. [unclear]".

Front End Processing Staff
Information Services Branch
Information Technology & Resources Management Division



+1 919.379.2510
+1 919.467.5923
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560
www.nufarm.com

Via Courier Delivery

October 1, 2014

Ms. Venus Eagle, PM #1
USEPA (7504P)
Document Processing Desk (AMEND)
Room S4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

Contains Confidential Business Information

**Subject: Nuprid 4F Insecticide, EPA Reg. No. 228-528
Application for Alternate CSF #9**

Dear Ms. Eagle:

Enclosed please find application for alternate CSF #9 for the above-referenced product. The purpose of this alternate CSF is as follows:


- Incorporate additional producers (CSF box 2);
- Incorporate additional suppliers (CSF boxes 10, 11, 12);
- Increase the amount of [REDACTED]
[REDACTED] to attain 100%.

Enclosed in support of this Application are the following:


- Application for Pesticide Registration - AMEND;
- One copy of Basic CSF; and
- Two copies of proposed, Alt. CSF #9.

Should you have any questions or require additional information, please contact me at (919) 379-2528 or by email at carrie.tackema@us.nufarm.com.

Sincerely,


Carrie M. Tackema
Regulatory Manager

Enclosure(s)



25



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☒ Amendment
☐ Other:

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 228-528	2. EPA Product Manager Venus Eagle	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Nuprid 4F Insecticide	PM# 1	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 11901 S. Austin Avenue Alsip, IL 60803 <i>Please send all correspondence to "contact point" listed below</i>		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input checked="" type="checkbox"/> Amendment – Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

Application for Alternate CSF #9

CONTACT: Carrie M. Tackema (919) 379-2528 or carrie.tackema@us.nufarm.com
Nufarm Americas, Inc.
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other	
*Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 16, 32, 64 fl. oz.; 1, 2.5, 220, 250 gal.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)					
Name Carrie M. Tackema		Title Regulatory Manager		Telephone No. (Include Area Code) (919) 379-2528	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature 		3. Title Regulatory Manager			
4. Typed Name Carrie M. Tackema		4. Date October 1, 2014			

DATA EXTRACTION REQUEST

Reg # 228-528 Decision # 492708

Description: Amendment

Material Sent:

- ☒ Electronic Label/Letter Dated 7/9/14
(See PPLS for electronic file)
- ☐ Stamped Label Dated _____ (See jacket)
- ☐ Notification Dated _____ (See jacket)
- ☐ New CSF(s) Dated _____ (See jacket)
- ☐ Other: _____

File this coversheet and attached materials in the jacket. It must be well organized and clipped together, NOT STAPLED. Then give the jacket with the coversheet and materials to staff in the Information Services Center (ISC) (Room S-4900). If a jacket is full or only available as an image, please file materials in a new jacket and bring it down to the (ISC). For further information please call 703-605-0716.

Reviewer: Jennifer Urbanski

Division: RD

Phone: 703-347-0156

Date: 7/11/14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 9, 2014

Carrie Tackema
Nufarm Americas, Inc.
11901 S. Austin Avenue
Alsip, IL 60803

Subject: Amended label removing application restriction to poultry houses within 7 days
of birds being present
Product Name: Nuprid 4F Insecticide
EPA Reg. No. 228-528
Submission dated June 16, 2014
Decision Number 492708

Dear Ms. Tackema:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one copy of your final printed labeling before you release the product for shipment. If you have any questions regarding this label, please contact Dr. Jennifer Urbanski at 703-347-0156 or urbanski.jennifer@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Venus Eagle" followed by a long horizontal flourish.

Venus Eagle, Product Manager 01
Insecticide-Rodenticide Branch
Registration Division
Office of Pesticide Programs

Enclosure- Stamped Label

Note: This Master label contains 2 Sub-labels which bear directions for use in Commercial Agriculture; and in Nursery, Greenhouse and Landscape Ornamentals

ACCEPTED

07/09/2014

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 228-528

NUPRID[®] 4F

INSECTICIDE

SUB-LABEL A: COMMERCIAL AGRICULTURE

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE ORNAMENTALS

ACTIVE INGREDIENT:

Imidacloprid, 1 -[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine..... 40.4%

OTHER INGREDIENTS:..... 59.6%

TOTAL..... 100.0%

EPA REG. NO. 228-528

EPA EST. NO. _____

MANUFACTURED FOR
NUFARM AMERICAS INC.
11901 SOUTH AUSTIN AVENUE
ALSIP, IL 60803



[Nufarm: Grow a better tomorrow]

[Grow a better tomorrow]

NUPRID[®] 4F

INSECTICIDE

[alternate brand names]

NUPRID[®] 4F MAX Insecticide

A SYSTEMIC AND FOLIAR INSECTICIDE FOR USE ON LISTED FIELD CROPS including COTTON, TOBACCO, POTATO, SOYBEANS and PEANUTS; IN CITRUS, TREE NUT, and FRUIT ORCHARDS; ON FIELD and GREENHOUSE VEGETABLES; ON BERRY, BUSH and VINE CROPS; and ON OTHER LISTED CROPS

ACTIVE INGREDIENT:

Imidacloprid, 1 -[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine..... 40.4%

OTHER INGREDIENTS:..... 59.6%**TOTAL**..... 100.0%

Contains 4 pounds of imidacloprid per gallon.

KEEP OUT OF REACH OF CHILDREN**CAUTION****SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-528

EPA EST. NO. _____

MANUFACTURED FOR
NUFARM AMERICAS INC.
11901 SOUTH AUSTIN AVENUE
ALSIP, IL 60803



NET CONTENTS _____ GALS. (_____ Liters)

[Nufarm: Grow a better tomorrow]

[Grow a better tomorrow]

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid contact with eyes or clothing. Wear protective eye wear. Wear long sleeved shirt and long pants, socks, shoes, and chemical resistant gloves made of waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

FIRST AID

IF INHALED	<ul style="list-style-type: none"> • Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User must:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. **DO NOT** apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates. This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note to Reviewer: the two statements in brackets below may be used as they relate to *Tilia* species:

[**DO NOT** apply this product, by any application method, to linden, basswood or other *Tilia* species in the State of Oregon,]

[**DO NOT** apply this product, by any application method, to linden, basswood or other *Tilia* species.]

See individual crops for specific pollinator protection application restrictions. If none exist under the specific crop, for foliar applications, follow these directions for use for crops that are contracted to have pollinator services or for food/feed and commercially grown ornamentals that are attractive to pollinators:

1. FOR CROPS UNDER CONTRACTED POLLINATION SERVICES



Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless the following condition has been met:

If an application must be made when managed bees are at the treatment site, the beekeeper providing the pollination services must be notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

2. FOR FOOD CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS



Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

SUB-LABEL A: COMMERCIAL AGRICULTURE

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

PRODUCT INFORMATION

Thorough uniform coverage is necessary to achieve insect control. A spray adjuvant may be used to improve coverage. This product may not knockdown established and heavy insect populations with a single application. Two applications may be required to achieve control; retreat if needed and as directed on this label. This product may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.

Applying this product to crops grown for production of true seed intended for private or commercial planting is not permitted unless allowed under state approved 24(c) labeling. Additional information on this product uses for listed crops and other questions may be obtained from the Cooperative Extension Service, PCA's, consultants or your local Nufarm representative.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product must conform to resistance management strategies established for your use area.

This product contains a Group 4A insecticide called imidacloprid. Insect biotypes with acquired or inherent tolerance to group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control of targeted species. This may eventually result in partial or total loss of control of those species by this product and other Group 4A products.

The active ingredient in this product is a member of neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of this product and/or other Group 4A products having the same or similar mode of action.

Following a neonicotinoid block of treatments, Nufarm strongly encourages the rotation to a block of applications with effective products of a different mode of action before using additional applications of neonicotinoid products. Use of a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect's pest's ability to develop resistance to this class of chemistry.

Foliar applications of this product or other Group 4A products from the neonicotinoid chemical class must not be used on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Intruder, Impulse, Leverage, Pasada, Provado, Trimax Pro and Venom. Other 4A Group, neonicotinoid products used as soil/seed treatment include: Admire Pro, Advise, Alias, Belay, Clutch, Couraze, Cruiser, Gaucho, Macho, Macho Max, Platinum, Venom and Widow. Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

PRODUCT USE INSTRUCTIONS

OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

SPRAY DRIFT MANAGEMENT

The responsibility of avoiding spray drift is with the applicator. The applicator must consider weather related factors and the interaction of application equipment when making application decisions.

Mixing and Loading Requirements

The use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well head, sinkholes or field drains.

Aerial Applications

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size

The droplet size is an important factor and can influence drift. Small droplets (<150 – 200 microns) drift to a greater extent than large droplets. Applications typically should be made to deliver the largest droplet range that provides adequate control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. **DO NOT** apply when winds are greater than 15 mph and avoid gusty and windless conditions.

Restrictions During Temperature Inversions

DO NOT make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions typically restrict vertical air mixing, which then could cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions typically are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices must be followed.

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- **DO NOT** allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

No-Spray Zone Requirements for Foliar Applications

DO NOT apply by ground within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

RUNOFF MANAGEMENT

DO NOT cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff must be employed.

ENDANGERED SPECIES NOTICE

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

SUB-LABEL A: COMMERCIAL AGRICULTURE

ROTATIONAL CROPS

Crops which are listed on imidacloprid labels or crops that have existing tolerances for imidacloprid may be planted in treated areas as soon as practical after the last imidacloprid application. Crops that are not found on an imidacloprid label, or crops that do not have existing tolerances for imidacloprid, may not be planted in treated areas for 12-months after the last application. Refer to the table below for plantback intervals for different crops. Note that if cover crops are planted any time after an application of this product, those crops may not be grazed or harvested for food or feed.

ROTATIONAL CROPS - PLANT-BACK INTERVALS
IMMEDIATE PLANT-BACK: All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop & sweet), rapeseed, sorghum, sugar beet and wheat.
30-DAY PLANT-BACK: Cereals (including buckwheat, millet, oats, rice, rye, and triticale), soybeans and safflower
10-MONTH PLANT-BACK: Onions and bulb vegetables
12-MONTH PLANT-BACK: All other crops

APPLICATION INSTRUCTIONS

This product can be applied as a foliar spray, or as a soil treatment (see Crop Specific Restrictions and Limitations). Thorough uniform coverage is necessary to achieve insect control. Use adequate spray volumes, properly calibrated application equipment, and an adjuvant to improve coverage. Failure to provide adequate coverage and retention of this product on leaves and fruit may result in loss of insect control or delay in onset of activity.

This product may not knockdown established and heavy insect populations with a single application. Two applications may be required to achieve control. Scout fields and retreat if needed.

This product may be tank mixed with other insecticides as recommended for knockdown of pests or for improved control of other pests.

Apply this product with properly calibrated ground or aerial application equipment.

Minimum spray volumes unless otherwise specified on the Crop Specific Restrictions and Limitations section are 10 gallons/Acre by ground applications and 5 gallons/Acre through aerial equipment.

This product may also be applied by overhead chemigation (see **CHEMIGATION APPLICATION** section below) if allowed in crop specific recommended application section.

Apply specified rate per acre as a foliar spray as pest populations begin to build. **DO NOT** apply more than 0.5 lbs. active ingredient per acre, per year, regardless of formulation or method of application, unless specified within the **Crop Specific Restrictions and Limitations** for a given crop.

Mix Preparation

To prepare the application mixture:

1. Fill the spray tank with a portion of the required amount of water and begin agitation.
2. Add the specified amount of NUPRID® 4F Insecticide.
3. Fill the tank with the remaining water needed. Maintain sufficient agitation during mixing and application.

NOTE: This product may also be used with other pesticides and/or fertilizer solutions; refer to the Tank Mix and Compatibility Note below. When tank mixtures of this product and other pesticides are involved, prepare the tank mixture as specified above and follow the suggested Mixing Order below.

Tank Mixes

Unless otherwise prohibited on this label or the label of an intended tank mix product, this product may be applied in combination with any pesticide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. **IMPORTANT: PESTICIDE TANK MIXES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS**

SUB-LABEL A: COMMERCIAL AGRICULTURE

LABEL OR IN MANUFACTURER'S SUPPLEMENTAL OR 24(C) LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

Compatibility

Before full-scale mixing of this product with other pesticides, fertilizers, secondary plant nutrients, adjuvants, surfactants or oils, you must determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent. **IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL OR 24(c) LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.**

Mixing Order

When pesticide mixtures are needed, add wettable powders first, this product or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added and do not add an additional component until the previous is thoroughly mixed. A fertilizer/pesticide compatibility agent may be needed if a fertilizer solution is to be added to the mixture. Be sure to maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

Further information on Tank Mixes is available from your local Nufarm representative.

CHEMIGATION APPLICATION

Types of Irrigation Systems

Chemigation applications of this product may be made to crops through overhead sprinkler chemigation systems if specified in crop-specific recommendation sections. **DO NOT** apply this product through any other type of irrigation system.

Water Volume

Chemigation applications of this product must be made as concentrated as possible. Retention of this product on target site of insect infestation is necessary for optimum activity. Chemigation of this product in water volumes exceeding 0.10 inches/Acre are not recommended.

Uniform Water Distribution and System Calibration

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift

DO NOT apply when the wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or normally shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems

SUB-LABEL A: COMMERCIAL AGRICULTURE

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

CROP-SPECIFIC RESTRICTIONS AND LIMITATIONS**FIELD CROPS**

COTTON		
SOIL APPLICATIONS		
Pests	Fluid ounces/1,000 row-feet	Fluid ounces/Acre
For control of: Cotton aphid Plant bugs Thrips Whiteflies	0.65	8.5 – 10.6 (Depending on row-spacing)
Application Methods		
Apply specified dosage by one of the following methods: 1. In-furrow or narrow band spray. When applying as an in-furrow spray, direct application on or below the seed at planting; OR 2. Narrow band application below the eventual seed bed row in a bedding operation 7 or fewer days before planting; OR 3. Chemigation into root zone through low-pressure drip or trickle irrigation equipment.		
Restrictions		
Maximum amount of product allowed per year: 10.6 fluid ounces/Acre (0.33 lb AI/Acre) DO NOT apply more than 0.5 lbs. of active ingredient per acre per year of NUPRID® 4F Insecticide, Provado®, Trimax® or Leverage®, including seed treatment as Gaucho®, soil and foliar uses. DO NOT graze treated fields after any application of this product. See Resistance Management section of this label.		
FOLIAR APPLICATIONS		
Pests	Fluid ounces/Acre	
For control of: Cotton aphid Cotton fleahopper Bandedwinged whitefly Plant bugs (excludes <i>Lygus hesperus</i>) Green stink bug Southern green stink bug Bollworm/Budworm (ovicidal effect)	1.0 – 2.0	
For suppression of: Lygus bugs (<i>Lygus hesperus</i>) Whiteflies (other than banded winged whitefly)	2.0	
Application Methods		
Apply through properly calibrated ground, aerial, or chemigation application equipment.		
Restrictions		
Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 7 days		

SUB-LABEL A: COMMERCIAL AGRICULTURE

Maximum amount allowed per year: **10.0 fluid ounces/Acre** (0.31 lb. AI/A)

DO NOT apply more than 0.5 lbs. of active ingredient per acre per year of NUPRID® 4F Insecticide, Provado®, Trimax® or Leverage®, including seed treatment as Gaucho®, soil and foliar uses.

DO NOT graze treated fields after any application of this product.

TANK MIX APPLICATIONS

Pests (in addition to those listed above)	NUPRID 4F Rate Fluid ounces/Acre	BIDRIN 8 Rate ¹ Fluid ounces/Acre
For early season control of: Thrips	1.0 – 1.5	1.6 – 3.2
For mid to late season control of: Plant bugs Stink bugs (including Brown stink bug) Grasshoppers Saltmarsh caterpillar Cotton leafperforator		4.0 – 8.0

Remarks

This product can be tank mixed with other pesticides and/or fertilizer solutions. When tank-mixing this product with other pesticides, prepare the tank mixture as specified above in the **Mix Preparation** instructions section. Follow the following general mixing order:

1. Add wettable powders first;
2. Add this product or other flowables second;
3. Add emulsifiable concentrates last.

Be sure to maintain agitation as each component is added and do not add an additional component until the previous component is thoroughly integrated into the mixture. If a fertilizer solution is added, a fertilizer-pesticide compatibility agent may be needed.

Restrictions (in addition to those listed above for NUPRID 4F foliar applications)

¹ Refer to the BIDRIN 8 product label for specific use instructions; observe all use restrictions and precautions that appear on the BIDRIN 8 label.

POTATO

SOIL APPLICATIONS

Pests	Fluid ounces/1,000 row-feet	Fluid ounces/Acre
For control of: Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid	0.45 – 0.65	6.5 – 10.0
For suppression of: Wireworms (with in-furrow spray at planting)		
For suppression of disease symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis (PLRV)		

Application Methods

Apply specified dosage in one of the following methods:

1. In-furrow spray during planting directed on seed pieces or seed potatoes; OR
2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil; OR
3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil; OR
4. Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting. For effective pest control or suppression, applications of this product must be placed below soil surface and in contact with seed piece or within root zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of this product may be made in a 2 to 4 inch band (width of planter shoe opening) and completely covered.

Restrictions

Maximum amount of product allowed per year: **10.0 fluid ounces/Acre** (0.31 lb AI/Acre)

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Colorado Potato beetle Flea beetles Flea hoppers Psyllids	1.5

Application Methods

SUB-LABEL A: COMMERCIAL AGRICULTURE

Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum amount allowed per year: **6.4 fluid ounces/Acre** (0.20 lb. AI/A)

SEED PIECE APPLICATIONS¹

Pests	Fluid ounces/100 lbs seed	Fluid ounces/Acre
For control of: Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid Wireworms (seed piece protection)	0.2 – 0.4	4.0 – 8.0
For suppression of disease symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis (PLRV)	0.4	8.0

Application Methods

Apply specified dosage as a diluted spray onto seed pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part this product. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after this product's application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed pieces as soon as possible after treating avoiding prolonged exposure of seed pieces treated with this product to sunlight and in accordance with the recommendation of your local Extension specialist. Consult your local Nufarm representative or crop protection product dealer for information relevant to your area.

Remarks

¹ Based on a seeding rate of 2000 lbs/Acre

Restrictions

Maximum amount of product allowed per year: **10.0 fluid ounces/Acre** (0.31 lb AI/Acre)

DO NOT use treated seed pieces for food, feed, or fodder.

DO NOT apply any subsequent applications of this product (in-furrow), Gaucho[®], Leverage[®] or Provado[®] following a seed-piece treatment of this product.

Apply only in areas that are equipped to remove spray mist or dust or with adequate ventilation.

SOYBEAN**FOLIAR APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: Aphids Bean leaf beetle Cucumber beetles / Rootworm adults Japanese beetle (adults) Leafhoppers Whiteflies	1.5

Application Methods

Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Minimum interval between applications: **7 days**

Maximum amount allowed per year: **3.65 fluid ounces/Acre** (0.13 lb. AI/A)

¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

TOBACCO**SOIL APPLICATIONS**

Pests	Fluid ounces/1,000 plants (as seedling tray drench)	Fluid ounces/1,000 plants (in-furrow or transplant-water)
For control of: Aphids	0.5	0.7

SUB-LABEL A: COMMERCIAL AGRICULTURE

Flea beetles		
For control of: Mole crickets Whiteflies Wireworms	0.7 – 1.4	0.9 – 1.4
For suppression of disease symptoms of: Tomato spotted wilt virus (TSWV)	1.4	1.4
Application Methods		
Apply specified dosage of this product in one of the following methods: 1. Uniform, broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting, followed immediately by overhead irrigation to wash this product from foliage into potting media. Failure to wash this product from foliage may result in a reduction in pest control. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.; OR 2. In-furrow spray or transplant-water drench during setting.; OR 3. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.		
Remarks		
Important Note: Proper tray drench applications of this product have been shown to be the most efficacious method of application. However, the specified rate of this product may be applied as a combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of this product into the plant and a delay in control.		
Restrictions		
Pre-Harvest Interval (PHI): 14 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb AI/Acre)		
FOLIAR APPLICATIONS		
Pests	Fluid ounces/Acre	
For control of: Aphids	0.8 – 1.6	
For control of: Flea beetles Japanese beetles	1.6	
Application Methods		
Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.		
Restrictions		
Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 7 days Maximum amount allowed per year: 9.0 fluid ounces/Acre (0.28 lb. AI/A)		

PEANUT	
SOIL APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	8.0 – 12.0
For suppression of: Thrips	12.0
Application Methods	
Apply as a: 1. Chemigation into root-zone through properly calibrated low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.; OR 2. In-furrow spray directed on or below seed.	
Remarks	
Applications of this product have been shown to increase the incidence of Tomato spotted wilt virus (TSWV), and possibly other tospoviruses, on multiple varieties of peanut. Prior to making product applications, contact the State, Cooperative This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling. Extension Service, or Nufarm representative, for recommendations to discuss the risk and benefits of imidacloprid applications.	
Restrictions	

SUB-LABEL A: COMMERCIAL AGRICULTURE

Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	1.4
Application Methods	
Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 5 days Maximum amount allowed per year: 4.2 fluid ounces/Acre (0.13 lb. AI/A) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	

VEGETABLE CROPS

CUCURBIT VEGETABLES: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i> including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of <i>Citrullus lanatus</i>).	
SOIL APPLICATIONS (FIELD)	
Pests	Fluid ounces/Acre
For control of: Aphids Cucumber beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	8.0 – 12.0
For suppression of disease symptoms of: Bacterial wilt (as vectored by various cucumber beetles) Leaf silvering resulting from whitefly feeding	12.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. In-furrow spray directed on or below seed; OR 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR 5. Post-seeding drench, transplant-water drench, or hill drench; OR 6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.	
Restrictions	
Pre-Harvest Interval (PHI): 21 days Maximum amount of product allowed per crop season: 12.0 fluid ounces/Acre (0.38 lb AI/Acre) DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling	
GREENHOUSE APPLICATIONS ¹	
Pests	Fluid ounces/1000 plants
For control of: Aphids Whiteflies	0.05
Application Method	
Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:	

SUB-LABEL A: COMMERCIAL AGRICULTURE

<ol style="list-style-type: none"> 1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash this product from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash this product from foliage may result in reduced pest control; OR 2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.
Remarks
<p>The application made in the planthouse is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.</p> <p>Important Note: Not all varieties of cucurbit vegetables have been tested for tolerance to this product applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.</p>
Restrictions
<p>Maximum number of greenhouse applications: 1</p> <p>Maximum amount of product allowed per greenhouse application: 0.05 fluid ounce (0.001568 lb AI)/1,000 plants.</p> <p>DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.</p> <p>¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.</p>

FRUITING VEGETABLES: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo.	
SOIL APPLICATIONS (FIELD)	
Pests	Fluid ounces/Acre
For control of: Aphids Colorado potato beetles Flea beetles Leafhoppers Thrips (foliage-feeding only) Whiteflies	Okra & Pepper 8.0 – 16.0
For suppression of disease symptoms of: Tomato mottle virus Tomato spotted wilt virus Tomato yellow leaf curl virus	Other Listed Crops 8.0 – 12.0
Application Methods	
Apply specified dosage of this product in one of the following methods: <ol style="list-style-type: none"> 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. In-furrow spray directed on or below seed; OR 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR 5. Post-seeding drench, transplant-water drench, or hill drench; OR 6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone. 	
Restrictions	
Pre-Harvest Interval (PHI): 21 days Maximum amount of product allowed on Okra and Pepper per crop season: 16.0 fluid ounces/Acre (0.5 lb AI/Acre). Maximum amount of product allowed on other listed fruiting vegetable crops per crop season: 12.0 fluid ounces/Acre (0.38 lb AI/Acre). DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS (FIELD)	
Pests	Fluid ounces/1000 plants
For control of: Aphids Colorado potato beetle Leaf beetles Whiteflies ¹	1.5 – 2.5
For control of: Pepper weevil (Pepper only) ²	2.5
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Remarks	
Applications of this product must be incorporated into a full-season program, where alternations of effective products from	

SUB-LABEL A: COMMERCIAL AGRICULTURE

multiple classes of chemistry and different modes of actions are utilized in a blocked or windowed approach. For additional information, please contact your Nufarm representative, Extension Specialist or crop advisor.

¹Higher specified rate within the rate range must be used when targeting adult whiteflies.

²For pepper weevil, apply specified dosage of this product by ground equipment only. Time applications prior to a damaging pest population becoming established. Good coverage of foliage and fruit is necessary for target pest control.

Restrictions

Pre-Harvest Interval (PHI): **0 days**

Minimum interval between applications: **5 days**

Maximum amount of product allowed per crop season: **7.7 fluid ounce (0.24 lb AI)/A**

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

GREENHOUSE APPLICATIONS¹

Pests	Fluid ounces/1000 plants
For control of: Aphids Whiteflies	0.05

Application Methods

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash this product from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash this product from foliage may result in reduced pest control; OR
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

Remarks

The application made in the planthouse is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.

Important Note: Not all varieties of cucurbit vegetables have been tested for tolerance to this product applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

Restrictions

Maximum number of greenhouse applications allowed: **1**

Maximum amount of product allowed per greenhouse application: **0.05 fluid ounce (0.001568 lb AI)/1,000 plants.**

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

¹This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

GREENHOUSE VEGETABLES: Mature Cucumber and Tomato plants in production greenhouses ONLY.

Pests	Fluid ounces/1000 plants
For control of: Aphids Whiteflies	0.7

Application Methods

Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. **DO NOT** apply to immature plants since phytotoxicity may occur.

Remarks

Make application when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (*Onius* sp.) can occur when this product is applied.

Many varieties of vegetables have been tested for tolerance to this product and show good safety. However, certain varieties may show more sensitivity to this product. Therefore, treat a few plants before treating the whole greenhouse.

Restrictions

Pre-Harvest Interval (PHI): **0 days**

Maximum number of applications per crop season: **1**

Maximum amount of product allowed per crop season: **0.7 fluid ounce (0.022 lb. AI)/1,000 plants.**

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

GLOBE ARTICHOKE

SOIL APPLICATION ¹	
Pests	Fluid ounces/1000 plants

SUB-LABEL A: COMMERCIAL AGRICULTURE

For control of: Aphids Leafhoppers	8.0 – 16.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. In-furrow spray directed on or below seed.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Maximum amount of product allowed per year: 16.0 fluid ounce/Acre (0.50 lb. AI/A) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATION	
Pests	Fluid ounces/1000 plants
For control of: Aphids Leafhoppers	1.6 – 4.0
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum Interval between applications: 14 days Maximum amount of product allowed per year: 16.0 fluid ounce/Acre (0.50 lb. AI/A)	

HERBS: Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

SOIL APPLICATIONS ¹ (FIELD)	
Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	8.0 – 12.0
For suppression of: Thrips (foliage-feeding only)	
Application Methods	
Apply specified dosage in one of the following methods: 1. In-furrow spray during planting directed on or below seed; OR 2. In-furrow spray or transplant-water drench during setting or transplanting; OR 3. Shanked-into or below eventual seed-line; OR 4. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	
Remarks	
Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat only small areas or numbers of plants and evaluate prior to full-scale use.	
Restrictions	
Pre-Harvest Interval (PHI): 14 days Maximum amount of product allowed per crop season: 12.0 fluid ounces/Acre (0.38 lb AI/Acre) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS (FIELD)	
Pests	Fluid ounces/Acre
For control of: Aphids Leaf beetles Leafhoppers Whiteflies	1.4
Application Methods	
Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Remarks	

SUB-LABEL A: COMMERCIAL AGRICULTURE

The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage and control.

Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat only small areas or numbers of plants and evaluate prior to full-scale use.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **5 days**

Maximum amount of product allowed per crop season: **4.2 fluid ounce (0.13 lb AI/A)**

HEAD and STEM BRASSICA VEGETABLES¹: Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gailon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, plus Turnip tops (leaves).

LEAFY GREENS VEGETABLES¹: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Raddicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), **Watercress²** (including upland).

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre (on 36 in. rows)
For control of: Aphids Leafhoppers Thrips (foliage-feeding only) Whiteflies	5.0 – 12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray directed on or below seed; OR
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR
5. Post-seeding drench, transplant-water drench, or hill drench; OR
6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre (0.38 lb AI/Acre)**

¹ **DO NOT** use on crops grown for seed unless permitted by state approved 24(c) labeling.

² For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following application. Applications must be made to fully leafed-up canopies only. **DO NOT** apply to native cress growing in streams or other bodies of water.

FOLIAR APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	1.5

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **5 days**

Maximum amount of product allowed per crop season: **7.7 fluid ounces/Acre (0.23 lb. AI/A)**

¹ **DO NOT** use on crops grown for seed unless permitted by state approved 24(c) labeling.

² For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following application. Applications must be made to fully leafed-up canopies only. **DO NOT** apply to native cress growing in streams or other bodies of water.

LEAFY PETIOLE VEGETABLES: Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, finocchio), Rhubarb, Swiss chard	
SOIL APPLICATIONS (FIELD)	
Pests	Fluid ounces/Acre (on 36 in. rows)
For control of: Aphids Leafhoppers Thrips (foliage-feeding only) Whiteflies	5.0 – 12.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. In-furrow spray directed on or below seed; OR 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR 5. Post-seeding drench, transplant-water drench, or hill drench; OR 6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.	
Restrictions	
Pre-Harvest Interval (PHI): 45 days Maximum amount of product allowed per crop season: 12.0 fluid ounces/Acre (0.38 lb AI/Acre) DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.	

LEGUME VEGETABLES (except Soybean, dry):	
Edible podded and Succulent shelled pea and Bean and Dried Shelled Pea and Bean including:	
Bean - <i>Lupinus</i> spp. (grain lupin, sweet lupin, white lupin, and white sweet lupin)	
Bean - <i>Phaseolus</i> spp. (field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)	
Bean - <i>Vigna</i> spp. (adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)	
Pea - <i>Pisum</i> spp. (dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)	
Other Beans and Peas - Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean.	
SOIL APPLICATIONS (FIELD)	
Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	8.0 – 12.0
For suppression of disease symptoms of: Bean common mosaic virus (BCMV) Bean golden mosaic virus (BGMV) Beet curly top hybrigeminivirus (BCTV)	
Application Methods	
Apply specified dosage of this product in one of the following methods:	
1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR	
2. In-furrow spray at planting directed on or below seed; OR	
3. In a narrow (2" or less) surface band over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours following application; OR	
4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; OR	
5. As a post-seeding drench, transplant-water drench, or hill drench.	
Restrictions	
Pre-Harvest Interval (PHI): 21 days	
Maximum amount of product allowed per crop season: 12.0 fluid ounces/Acre (0.38 lb AI/Acre)	
DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.	
FOLIAR APPLICATIONS (FIELD)	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	1.4

SUB-LABEL A: COMMERCIAL AGRICULTURE

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum amount of product allowed per crop season: **4.2 fluid ounce** (0.13 lb AI/A)

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

ROOT VEGETABLES: Beet (garden)[†], Burdock (edible)[†], Carrot[†], Celeriac[†], Chervil (turnip-rooted)[†], Chickory[†], Ginseng, Horseradish, Parsley (turnip-rooted)[†], Parsnip[†], Radish[†], Oriental radish (diakon)[†], Rutabaga[†], Salsify (black)[†], Salsify (oyster plant), Salsify (Spanish), Skirret, Turnip[†]

TUBEROUS and CORM VEGETABLES (except Potato): Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet)[†], Chayote (root), Chufa, Dasheen (taro)[†], Ginger, Leren, Sweetpotato, Tanier (cocoyam)[†], Turmeric, Yam bean (jicama, manioc pea), Yam (true)[†]

SOIL APPLICATIONS[†] (FIELD)

Pests	Fluid ounces/1,000 row ft.	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Thrips (foliage-feeding) Whiteflies	0.35 – 0.85	5.0 – 12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray (rate specified per 1,000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting; OR
3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting; OR
4. Side-dress not more than 0.6 fluid ounces/1000 row-feet no later than 45 days after planting.

Remarks

Important Note: The rate applied affects the length of control. Use higher specified rates within the rate range where infestations occur later in crop development, or where pest pressure is continuous. Rates of this product less than 0.7 fluid ounce/1,000 row-feet will not provide adequate residual pest control. Crops treated with this product grown on very high organic matter soils (muck) may also require additional pest management control.

[†] The tops or greens from these crops may be utilized for food or feed.

Restrictions

Pre-Harvest Interval (PHI) for Root Vegetables: **21 days**

Pre-Harvest Interval (PHI) for Tuberous and Corm Vegetables: **3 days** (leaves); **125 days** (corms)

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Maximum applications per crop season: **1**

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

[†] Soil application to Root Vegetables is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS[†] (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	1.4

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Remarks

[†] The tops or greens from these crops may be utilized for food or feed.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **5 days**

Maximum amount of product allowed per crop season:

Radish: **1.4 fluid ounce** (0.044 lb AI/A)

SUB-LABEL A: COMMERCIAL AGRICULTURE

All other listed crops: **4.2 fluid ounces/Acre** (0.13 lb. AI/A)

Maximum applications per crop season:

Radish: **1**

All other listed crops: **3**

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

SUGARBEET: (California only)

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	3.0 – 6.0
For suppression of disease symptoms of: Western yellows virus Beet curly top hybrigeminivirus (BCTV)	

Application Methods

Apply specified dosage of this product in the following method:

Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting.

Remarks

The low rate may be applied to aid establishment of stands in whitefly areas, or for early season control of the other pests listed.

Restrictions

Maximum amount of product allowed per year: **6.0 fluid ounces/Acre** (0.19 lb AI/Acre)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

NUPRID® 4F INSECTICIDE CONVERSION CHART FOR LINEAR APPLICATION ONLY

RATE fluid ounces/ Acre	RATE fluid ounces/1,000 row-feet Based on <u>average</u> row spacing (in inches):									
	10	15	20	25	30	34	36	38	40	45
5.0	0.10	0.14	0.19	0.24	0.29	0.33	0.34	0.36	0.38	0.43
5.5	0.11	0.16	0.21	0.26	0.32	0.36	0.38	0.40	0.42	0.47
6.0	0.11	0.17	0.23	0.29	0.34	0.39	0.41	0.44	0.46	0.52
6.5	0.12	0.19	0.25	0.31	0.37	0.42	0.45	0.47	0.50	0.56
7.0	0.13	0.20	0.27	0.33	0.40	0.46	0.48	0.51	0.54	0.60
7.5	0.14	0.22	0.29	0.36	0.43	0.49	0.52	0.55	0.57	0.65
8.0	0.15	0.23	0.31	0.38	0.46	0.52	0.55	0.58	0.61	0.69
8.5	0.16	0.24	0.33	0.41	0.49	0.55	0.59	0.62	0.65	0.73
9.0	0.17	0.26	0.34	0.43	0.52	0.59	0.62	0.65	0.69	0.77
9.5	0.18	0.27	0.36	0.45	0.55	0.62	0.65	0.69	0.73	0.82
10.0	0.19	0.29	0.38	0.48	0.57	0.65	0.69	0.73	0.77	0.86
10.5	0.20	0.30	0.40	0.50	0.60	0.68	0.72	0.76	0.80	0.90
11.0	0.21	0.32	0.42	0.53	0.63	0.72	0.76	0.80	0.84	0.95
11.5	0.22	0.33	0.44	0.55	0.66	0.75	0.79	0.84	0.88	0.99
12.0	0.23	0.34	0.46	0.57	0.69	0.78	0.83	0.87	0.92	1.03
12.5	0.24	0.36	0.48	0.60	0.72	0.81	0.86	0.91	0.96	1.08

SUB-LABEL A: COMMERCIAL AGRICULTURE

13.0		0.37	0.50	0.62	0.75	0.85	0.90	0.95	0.99	1.12
13.5		0.39	0.52	0.65	0.77	0.88	0.93	0.98	1.03	1.16
14.0		0.40	0.54	0.67	0.80	0.91	0.96	1.02	1.07	1.21

Important Note: Rate of this product applied affects the length of control and, to a considerable extent, the degree of control or effect. Row-spacing X rate combinations in shaded blocks may not provide adequate residual pest control and are not recommended for long-term, residual control. Use higher labeled rates where infestations may occur later in crop development or where pest pressure is continuous. Nufarm offers no warranty for use of this product at rates below 0.35 fluid ounce/1,000 row- feet (the Row-Spacing/Rate combinations that are shaded).

BERRY, BUSH and VINE CROPS

STRAWBERRY: annual and perennial varieties	
SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Whiteflies	12.0 – 16.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening; OR 2. As a plant material or plant hole treatment just prior to, or during transplanting; OR 3. As a pre-plant band spray over the row in a minimum of 20 gallons of water per acre, followed immediately by overhead irrigation to incorporate product into root zone. DO NOT use plastic or other mulch that limits movement of this product into root zone.	
Remarks	
The rate applied affects the length of control. Use higher specified rates within the rate range where infestations may occur later in crop development or where pest pressure is continuous.	
Restrictions	
Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per crop season: 16.0 fluid ounces/Acre (0.50 lb AI/Acre) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. DO NOT make both a soil and foliar application on the same crop in the same season. DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.	
SOIL APPLICATIONS (post-harvest use on perennial varieties)	
Pests	Fluid ounces/Acre
For control of: White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	8.0 – 12.0
Application Methods	
Apply a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply specified dosage of this product in one of the following methods: 1. As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre; OR 2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth must be equivalent to the width of the anticipated fruiting bed; OR 3. As a chemigation application with 600 to 1,000 gallons of water followed by 0.10 to 0.25 inch irrigation.	
Remarks	
All soil-surface applications must be followed by 0.25 inch of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate this product into egg-deposition zone may result in decreased activity of beetle grubs.	
Restrictions	
Maximum amount allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A)	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Spittlebugs	1.5

SUB-LABEL A: COMMERCIAL AGRICULTURE

Whiteflies
Application Methods
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.
Remarks
All soil-surface applications must be followed by 0.25 inch of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate this product into egg-deposition zone may result in decreased activity of beetle grubs.
Restrictions
Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum amount of product allowed per crop season: 4.6 fluid ounces/Acre (0.14 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. DO NOT make both a soil and foliar application on the same crop in the same season. DO NOT use on crops grown for seed unless allowed by state approved 24(c)labeling.

BUSHBERRY: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal	
SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Japanese beetle (adults, feeding on foliage) White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. 18-inch band on each side of the row followed with 0.25 inch of irrigation immediately after application.	
Remarks	
For grub control, apply this product to control 1st or 2nd (early) instar larvae. Application may be made post-bloom up to 7 days prior to harvest, or post-harvest until October 1st. For control of Japanese beetle larvae, make applications from June 1 to July 15. DO NOT apply during bloom. Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root zone will help protect berry plant roots from grub feeding. Apply this product to moist soil. If necessary, apply one hour of irrigation water immediately before application. To ensure maximum efficacy, 0.5 to 1 inch of irrigation water or rainfall must be applied or received within 24 hours of application of this product to facilitate movement into the soil and into the root zone.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb AI/Acre) DO NOT apply pre-bloom or during bloom or when bees are foraging.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters	1.2 – 1.6
For control of: Japanese beetles (adults) Thrips (foliage feeding)	2.4 – 3.2
For control of: Blueberry maggot	3.2
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 7 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) Maximum number of product applications per year: 5 Maximum application volume (water): Ground: 20.0 GPA; Air: 5.0 GPA DO NOT apply pre-bloom or during bloom or when bees are foraging.	

SUB-LABEL A: COMMERCIAL AGRICULTURE

CANEBERRY: Blackberry (<i>Rubus eubatus</i> , including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these). Raspberry (black and red, <i>Rubus occidentalis</i> , <i>Rubus strigosus</i> , <i>Rubus idaeus</i>).	
SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	8.0 – 16.0
For control of: Rednecked cane borer	12.0 – 16.0
For suppression of: Thrips (foliage-feeding only)	16.0
Application Methods	
Apply specified dosage in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. Basal, soil drench in a minimum of 500 gallons solution per acre.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days . Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb AI/Acre) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.	
FOLIAR APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers	3.2
For suppression of: Thrips (foliage-feeding only)	
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 7 days Maximum amount of product allowed per season: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are actively foraging.	
¹ This use is not permitted in CA unless otherwise directed by approved 24(c) labeling.	

CRANBERRY	
SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Rootgrubs (Scarab) Rootworms (Chrysomelid)	8.0 – 16.0
Application Methods	
Apply this product to moist soil. Apply specified dosage of this product in one of the following methods: 1. As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre; OR 2. As a chemigation application with 600 to 1,000 gallons water.	
Immediately upon application, this product must be incorporated into root zone by 0.1 to 0.3 inch water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.	
Make application post-bloom immediately after honeybees are removed. Application should target early instar larvae.	
Remarks	
Best control may be achieved when application is made post-bloom immediately after bees are removed. Target early instar larvae. This product has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of this product and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.	

SUB-LABEL A: COMMERCIAL AGRICULTURE

Restrictions

Pre-Harvest Interval (PHI): **30 days**
 Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)
DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

GRAPES: American bunch grape, Muscadine grape, and Vinifera grape.

SOIL APPLICATIONS

Pests	Fluid ounces/Acre
For control of: European fruit lecanium Leafhoppers/Sharpshooters Mealybugs <i>Phylloxera</i> spp. ¹	8.0 – 16.0
For suppression of: Grapeleaf skeletonizer Nematodes ²	16.0
For suppression of disease symptoms of : Pierce's disease	

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation; OR
3. Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation.

Remarks

Make application between bud-break and the pea-berry stage. A total of 14 fluid ounces/acre is required under the following conditions:

1. Where vigorous vine growth is expected
2. In warmer growing areas
3. Where mealybug and European fruit lecanium populations are expected to be heavy
4. Where vine populations exceed 600 per acre, or;
5. For suppression of nematodes

¹ Repeated and regular use of this product over multiple consecutive growing seasons controls existing *Phylloxera* infestations over time or prevents *Phylloxera* from becoming established.

²For suppression of nematodes, apply 14 fluid ounces in a single application or two 7-fluid ounce applications on a 30 to 45-day interval. Only make treatments by 1) chemigation into root zone through above ground low pressure drip, tickle, micro sprinkler or equivalent equipment or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root zone of the plant. Repeated and regular use of this product over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

Restrictions

Pre-Harvest Interval (PHI): **30 days**
 Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Leafhoppers/Sharpshooters Mealybugs	1.2 – 1.6
For control of: Grapeleaf skeletonizer	1.6

Application Methods

Apply specific dosage of this product using properly calibrated ground application equipment only. Apply as a broadcast or directed spray to infested areas ensuring thorough coverage.

Restrictions

Pre-Harvest Interval (PHI): **0 days**
 Minimum interval between applications: **14 days**
 Maximum amount of product allowed per year: **3.2 fluid ounces/Acre** (0.1 lb. AI/A)

HOPS:

SOIL APPLICATIONS¹

Pests	Fluid ounces/Acre
-------	-------------------

SUB-LABEL A: COMMERCIAL AGRICULTURE

For control of: Aphids	9.6
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation; OR 3. Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation.	
Restrictions	
Pre-Harvest Interval (PHI): 60 days Maximum amount of product allowed per year: 9.6 fluid ounces/Acre (0.3 lb AI/Acre) ¹ This use is not permitted in CA unless otherwise specified by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids	3.2
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 28 days Minimum interval between applications: 21 days Maximum amount of product allowed per year: 9.6 fluid ounces/Acre (0.3 lb. AI/A)	

COFFEE:

SOIL APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Leafminers	8.0 – 16.0
For suppression of: Scales	
Application Methods	
Apply specified dosage in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.; OR 2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation.; OR 3. Basal, soil drench in sufficient water to insure incorporation into the root zone followed by irrigation.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb AI/Acre) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. ¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	3.2
For suppression of: Scales	
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 7 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.	

CITRUS, TREE NUT and ORCHARD CROPS

CITRUS (containerized): Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

SOIL APPLICATIONS

Pests	mL/0.1ft ³ of container media
For control of: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies Citrus root weevil (larval complex) ¹	0.38-0.58
For suppression of: Citrus thrips (foliage-feeding only)	0.58

Application Methods

Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of this product per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, treatment must be made at planting prior to insect infestation. Retreat if necessary.

Application Restrictions

Pre-Harvest Interval (PHI): **0 day**
Maximum allowed per application: **0.58 mLs/0.1 ft³ container media**
Maximum allowed per crop season: **3.5 mLs/plant**
Do not apply pre-bloom or during bloom or when bees are foraging.

Remarks

¹For control of larvae of the citrus root weevil complex, make application prior to neonate larvae entering potting media. Utilize specified higher dosage for heavy infestations.

CITRUS: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

SOIL APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Termites (FL only) Whiteflies	8.0 – 16.0
For suppression of: Citrus nematode Thrips (foliage-feeding thrips only)	16
For suppression of disease symptoms of: Citrus tristeza virus (CTV) through vector control Citrus yellows	

Application Methods

Apply specified dosage of this product in one of the following methods:

- Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. Apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications of this product. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move this product into root zone. Allow 24 hours before initiating subsequent irrigations; OR
- Soil surface band spray on both sides of the tree. Bands must overlap at the tree base to create a continuous band within the

SUB-LABEL A: COMMERCIAL AGRICULTURE

<p>drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root zone. This method is suitable for very coarse soils with 0.75% organic matter or less; OR</p> <p>3. Drench to base of tree not exceeding one quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only for trees up to 8 feet tall; OR</p> <p>4. For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk; OR</p> <p>5. For suppression of citrus nematode, apply specific dosage through low pressure chemigation or soil surface spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of this product over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.</p>		
Restrictions		
<p>Pre-Harvest Interval (PHI): 0 days</p> <p>Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A)</p> <p>DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.</p>		
FOLIAR APPLICATIONS		
Pests	Fluid ounces/100 gallons	Fluid ounces/Acre
For control of: Aphids Asian citrus psyllid Black fly Leafhoppers/Sharpshooters Leafminers Mealybugs Scales ¹ Whiteflies	1.4 – 2.0 (dilute application)	4.0 – 8.0 (depending on tree size, target pest, and infestation pressure)
For suppression of: Thrips (foliage-feeding thrips only)	2.0	8.0
Application Methods		
<p>Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply through properly calibrated ground or aerial equipment.</p>		
Remarks		
<p>Aerial application of this product may result in slower activity and reduced control compared to ground application. Where higher rate applications are appropriate, increase the spray solution concentration to apply an equivalent rate per acre to that applied in the diluted application. The 8.0 fluid ounce/Acre rate is based on full sized trees. This rate may be reduced proportionally for smaller trees.</p> <p>¹Scales - time applications to the crawler stage. Treat each generation.</p>		
Restrictions		
<p>Pre-Harvest Interval (PHI): 0 days</p> <p>Minimum interval between sprays: 10 days</p> <p>Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A)</p> <p>DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.</p>		

POME FRUIT: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince		
SOIL APPLICATIONS		
Pests	Fluid ounces/Acre	
For control of: Aphids (including Wooly apple aphid) Leafhoppers	8.0 – 12.0	
Application Methods		
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.		
Restrictions		
Pre-Harvest Interval (PHI): 21 days Maximum amount of product allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.		
FOLIAR APPLICATIONS		
Pests	Fluid ounces/100 gallons	Fluid ounces/Acre
For control of: Leafhoppers	0.4 – 0.8	1.6 – 3.2
For control of: Aphids (except Woolly apple aphid) Apple maggot Leafminers	0.8	3.2

SUB-LABEL A: COMMERCIAL AGRICULTURE

San Jose scale		
For use on Pears Only to control: Mealybugs Pear psylla	2.0	8.0
Application Methods		
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.		
Remarks		
Combine applications targeting apple maggot with manufacturer's specified rate of a sticker.		
Restrictions		
Pre-Harvest Interval (PHI): 7 days Minimum interval between sprays: 10 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.		

POMEGRANATE	
SOIL APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Whiteflies	8.0 – 16.0
Application Methods	
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 0 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Whiteflies	3.2
For suppression of: Scales	
Application Methods	
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between sprays: 7 days Maximum amount of product allowed per year: 9.6 fluid ounces/Acre (0.3 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.	

STONE FRUIT: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)	
PRE-PLANT ROOT DIP APPLICATIONS	
Pests	Fluid ounces/10 gallons root dip solution
For control of: Black peach aphid (infesting roots)	1.0
Application Methods	
Mix this product at a rate of 1.0 fluid ounce per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in this product's solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.	
SOIL APPLICATIONS	

SUB-LABEL A: COMMERCIAL AGRICULTURE

Pests		Fluid ounces/Acre	
For control of: Aphids (including Woolly apple aphid) Leafhoppers		8.0 – 12.0	
Application Methods			
Apply specified dosage of this product in the following method: Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.			
Restrictions			
Pre-Harvest Interval (PHI): 21 days Maximum amount of product allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.			
FOLIAR APPLICATIONS			
Pests		Fluid ounces/100 gallons	Fluid ounces/Acre
For control of: Aphids Green June beetle Japanese beetle Leafhoppers/Sharpshooters Plant bugs Rose chafer San Jose scale		0.8	1.6 – 3.2
For control of: Cherry fruit fly (maggot of Eastern & Western)			2.4 – 3.2
For suppression of: Plum curculio Stinkbugs			3.2
Application Methods			
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.			
Restrictions			
Apricot, Nectarine, Peach: Pre-Harvest Interval (PHI): 0 days Minimum interval between applications: 7 days Maximum amount of product allowed per crop season: 9.6 fluid ounces/Acre (0.30 lb. AI/A) Minimum application volume (water): Ground: 50 GPA; Air: 25 GPA DO NOT apply during bloom or within 10 days prior to bloom or when bees are actively foraging.			
Cherry, Plum, Plumcot, Prune: Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb. AI/A) Minimum application volume (water): Ground: 50 GPA; Air: 25 GPA DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.			

TREE NUTS except Almonds: Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)	
SOIL APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Mealybugs Spittlebugs Termites Two-lined spittlebugs Whiteflies	8.0 – 16.0
For suppression of: Thrips (foliage-feeding only)	16.0
For suppression of disease symptoms of: Pecan scab (from reduction in honeydew deposition)	
Application Methods	
Apply specified dosage prior to or at onset of pest infestation in one of the following methods:	
1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. Pre-wet soil prior to applications of this product and allow soil to dry following application and prior to subsequent irrigation: OR	

SUB-LABEL A: COMMERCIAL AGRICULTURE

<p>2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site; OR</p> <p>3. Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Apply this product in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area must follow within 48 hours to promote uptake by root system; OR</p> <p>4. For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18 to 24 inches. Allow soil to dry following treatment and prior to applying any irrigation.</p>	
Remarks	
Use higher specified rates within the rate range when applied by shank or subsurface side-dress, used on larger trees, soils with high clay content, for high plant populations, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.	
Restrictions	
<p>DO NOT apply in Almonds</p> <p>Pre-Harvest Interval (PHI): 7 days</p> <p>Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb. AI/A)</p> <p>DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.</p> <p>¹Soil Application in Tree Nut orchards is not permitted in California unless otherwise directed by state approved 24(c) labeling.</p>	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids (except Black pecan aphid) Leafhoppers/Sharpshooters Phylloxera spp. (leaf infestations) Spittlebugs Whiteflies	1.4 – 2.8
For control of: Black pecan aphid Mealybugs San Jose scale ¹	3.2
Application Methods	
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.	
Remarks	
¹ Applications for control of San Jose scale must be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.	
Restrictions	
<p>DO NOT apply on Almonds</p> <p>Pre-Harvest Interval (PHI): 7 days</p> <p>Minimum interval between applications: 6 days</p> <p>Maximum amount of product allowed per year: 11.5 fluid ounces/Acre (0.36 lb. AI/A)</p> <p>Minimum application volume (water): Ground: 50 GPA; Air: 25 GPA</p> <p>DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.</p>	

BANANA and PLANTAIN

SOIL APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers	8.0 – 16.0
For suppression of: Scales	
Application Methods	
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 0 days	
Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb. AI/A)	
¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre

SUB-LABEL A: COMMERCIAL AGRICULTURE

For control of: Aphids Leafhoppers Thrips	3.2
Application Methods	
Apply specified dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.	
Remarks	
Aerial application of this product may result in slower activity and reduced control relative to results from ground application. Addition of an organosilicone adjuvant at a rate not to exceed 2.0 fluid ounces/100 gallons finished spray solution may improve coverage and pest control.	
Restrictions	
Pre-Harvest Interval (PHI): 0 days Minimum interval between applications: 14 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A)	

TROPICAL FRUIT: Acerola, Atemoya ¹ , Avocado, Biriba ¹ , Black sapote, Canistel, Cherimoya ¹ , Custard apple ¹ , Feijoa, Jaboticaba, Guava, Llama ¹ , Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop ¹ , Spanish lime, Star apple, Starfruit, Sugar apple ¹ , Wax jambu.	
SOIL APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Avocado lacebug Leafhoppers Whiteflies	12.0 – 16.0
For suppression of: Scales Thrips (foliage-feeding thrips only)	16.0
Application Methods	
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 6 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb. AI/A) ¹ Soil application use on noted crops is not permitted in California unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Mealybugs Thrips Whiteflies	3.2
For suppression of: Thrips	
Application Methods	
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.	
Remarks	
Ground applications of this product are more effective than aerial applications.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A)	

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

OTHER CROPS**Christmas Trees****SOIL APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: White grub complex (e.g., grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0

Application Methods

Soil incorporation and movement of this product to the root zone is required for activity. This product can be incorporated most readily when applied to moist soil. Apply specified dosage in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25 to 1 inch of irrigation within 12 hours after application.

Remarks

Apply this product during adult flight activity, or up to mid-July, when first instar larvae are present.

Restrictions

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb. AI/A)

¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Adelgids Sawflies	1.6 – 3.2

Application Methods

Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.

Remarks

Ground applications of this product are more effective than aerial applications.

For gall-forming adelgids, time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. Once galls form spraying this product is ineffective.

Restrictions

Minimum interval between applications: **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

Poplar/Cottonwood: (including members of the genus *Populus* grown for pulp or timber)**SOIL APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: Aphids Cottonwood leaf beetle	8.0 – 16.0
For suppression of: <i>Phylloxera popularia</i>	

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation through low-pressure drip irrigation; OR
2. For narrow row, cutting orchards/nurseries used for plant propagation, shank into root zone followed by adequate irrigation to promote uptake. Adequate irrigation depends on soil moisture level at application. Under dry conditions 0.25 inch/acre is recommended.

Remarks

For Cottonwood leaf beetle, protection against damage will occur when application is made early-season, when beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake. For *Phylloxera*, apply early in the year, from break of dormancy through May.

Restrictions

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb. AI/A)

¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.

CUTTING/WHIP APPLICATIONS¹

SUB-LABEL A: COMMERCIAL AGRICULTURE

Pests	Soaking Solution Fluid ounces needed per 100 gallons
For control of: Cottonwood leaf beetle	6.65 to 13.3 (unhydrated cuttings/whips) 13.3 to 20.0 (partially hydrated cuttings/whips)
For suppression of: Aphids <i>Phylloxera popularia</i>	13.3 (unhydrated cuttings/whips) 20.0 (partially hydrated cuttings/whips)
Application Methods	
Apply this product in one of the following cuttings/whips soaking methods: 1. For freshly cut (hydrated) cuttings/whips, soak plant material in specified solution concentration for 24 hours prior to cold storage. After removal from cold storage, plant as needed; OR 2. For previously hydrated cuttings/whips removed from cold storage, allow plant material to reach room temperature and soak in specified solution concentration for 24 hours prior to planting. Take proper care in disposal of any residual soaking solution. Apply solution to existing trees or other registered crops as long as all product label precautions and restrictions are observed.	
Remarks	
The moisture content prior to application of the cuttings/whips, the solution concentration and the length of soaking interval interact to affect the amount of product absorbed into plant material. For a constant soaking interval of 24 hours, dry cuttings/whips absorb a higher quantity of solution and require a lower concentration. Conversely, more hydrated cuttings/whips absorb less solution and require a higher concentration. Soaking of cuttings/whips must occur in a covered container in absence of UV light. Not all <i>Populus</i> sp. clones/varieties/hybrids have been tested for crop safety. Without specific knowledge about a particular <i>Populus</i> sp. clone/variety/hybrid, a small number of cuttings/whips of each must be treated and evaluated prior to commercial use.	
Restrictions	
Maximum amount of product allowed at plant per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) ¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leaf beetles	1.6 – 3.2
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Remarks	
Ground application of this product is more effective than aerial application for these crops.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 10 days Maximum amount of product allowed per season: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or with 10 days prior to bloom or when honeybees are foraging. ¹ Use as a foliar application to Poplar/Cottonwood is not permitted in California unless otherwise directed by state approved 24(c) labeling.	

COMMERCIAL POULTRY FACILITIES

POULTRY HOUSING STRUCTURES	
Pests	Fluid ounces / 1,000 ft ²
For control of: Darkling beetles Hide beetles (Dermestids)	3.0 (90 ml)
Application Methods	
Apply between flocks after de-caking and sanitation procedures have been completed. Apply as a spot, crack and crevice, or surface spray on floors, walls, and support beams of structure. Apply using a minimum of 1/2 to 2 gallons of spray mixture per 1000 square feet. To prepare the spray mixture, fill the spray tank with 1/2 the required amount of water, then add the specified amount of product. Add the remaining water while agitating or mixing. Maintain constant agitation while applying.	
Apply spray mixture to the entire footing including 1 foot up the wall above the footing, and in 3 to 4 foot wide bands directly beneath all feed lines. The areas beneath the feed lines typically harbor large numbers of adult and larval stages of the target pest when an infestation occurs. Measure these areas to determine the appropriate amount of spray mixture to apply. For structures that are prone to large infestations, treat the footings including 1 foot up the wall and the entire floor area of the structure.	

SUB-LABEL A: COMMERCIAL AGRICULTURE

Cracks and crevice areas also are prone to large infestations of the target pest. Apply as a crack and crevice treatment around wall insulation or other areas that may harbor the target pest. If structures have supporting beams, treat the floor with a 1 foot band around each beam and apply 2 feet up the beam.

For structures prone to extreme infestation, treat the entire structure with a broadcast application. Apply 3.0 fluid ounces in 2 gallons of water per 1000 square feet of surface. Apply as a broadcast spray to areas where litter has accumulated (floor, under feed and water lines, lower sections of walls, corners).

Remarks

In order to avoid problems with pest resistance to imidacloprid, rotate to an insecticide with a different mode of action every 2-3 flocks. Rotate between 3 different insecticide mode of action classes labeled for control of target pests during a calendar year.

Restrictions

DO NOT apply when birds are present.

DO NOT allow food or feed to be contacted by the spray.

Remove feed and water from the treatment area before applying.

When treating the perimeter, do not allow this product to contact plants in bloom if bees are foraging the treatment area.

Pests	Fluid ounces / Gallon
For control of: Nuisance ants	0.125 – 0.25 (3/4 – 1.5 TSP)

Application Methods

Apply as a crack and crevice or wall void treatment inside structures. Apply to cracks, crevices, drilled holes, onto walls, around entry points such as doors, windows, vents, eaves, soffits, and utility access openings. If nests are present in voids, apply into the void if possible. Apply evenly to treatment surfaces but not to the point of runoff. Apply to areas around the exterior of the structure where ants may be present (soil, turf, ornamental shrubs and plantings, and groundcover in close proximity to or touching the structure). For above-ground nests, such as in wood posts, decks, or fences, or in trees, spray into holes/openings where ants are traveling and on the wood surface.

Restrictions

DO NOT use for control of native or imported fire ants, harvester ants or pharaoh ants.

Keep people and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL [HANDLING]:

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure

SUB-LABEL A: COMMERCIAL AGRICULTURE

rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable containers larger than 5 gallons]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[Refillable containers for return to Nufarm]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

SUB-LABEL A: COMMERCIAL AGRICULTURE

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

NUPRID is a registered trademark of Nufarm Americas Inc.

All other trademarks that appear on this label which are not owned by Nufarm Americas Inc. or its subsidiaries are the property of their respective owners.

(RV070914)

GROUP **4A** INSECTICIDE

NUPRID[®] 4F

INSECTICIDE

A SYSTEMIC and FOLIAR INSECTICIDE FOR USE on LISTED ORNAMENTALS, FRUIT and NUT TREES, and VEGETABLE PLANTS grown in LAWN and LANDSCAPE AREAS, in GREENHOUSES, NURSERIES, and INTERIOR PLANTSCAPES.

ACTIVE INGREDIENT:

Imidacloprid, 1 -[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine..... 40.4%

OTHER INGREDIENTS: 59.6%

TOTAL:..... 100.0%

Contains 4 pounds of imidacloprid per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-528
EPA EST. NO. _____

MANUFACTURED FOR
NUFARM AMERICAS INC.
11901 SOUTH AUSTIN AVENUE
ALSIP, IL 60803



NET CONTENTS _____ **GALS.** (_____ **Liters**)
[Nufarm: Grow a better tomorrow]
[Grow a better tomorrow]

000228-00528.20140709.draft

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid contact with eyes or clothing. Wear protective eye wear. Wear long sleeved shirt and long pants, socks, shoes, and chemical resistant gloves made of waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

PERSONAL PROTECTIVE EQUIPMENT (PPE)**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS**User must:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF INHALED	<ul style="list-style-type: none"> • Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. **DO NOT** apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates. This chemical demonstrates the properties and characteristics associated with

chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See individual sites for specific pollinator protection application restrictions. If none exist under the specific site, for foliar applications, follow these application directions for use for food/feed crops that are commercially grown ornamentals that are attractive to pollinators and for non-agricultural uses:

1. FOR FOOD CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS


Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

2. NON-AGRICULTURAL USE SITES


Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Shake well before using.

DO NOT apply when wind speed favors drift beyond the area intended for treatment.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE () AMENTALS

- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

- Keep children and pets off treated areas until sprays have dried.

PRODUCT INFORMATION

Thorough uniform coverage is necessary to achieve target pest control. A spray adjuvant may be used to improve coverage. This product may not knockdown established and heavy insect populations with a single application. Two applications may be required to achieve control; retreat if needed and as directed on this label. This product may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.

Apply this product as a broadcast or directed spray application. Time applications to begin as target pest populations begin to build. Ensure that the treated area receives a thorough, uniform coverage of the spray solution. To improve coverage, a spray adjuvant may be added to the NUPRID® 4F Insecticide tank mix.

Apply using either ground equipment in a minimum spray volume of 10 gallons per acre or with aerial equipment in a minimum spray volume of 5 gallons of water per acre. Use adequate spray volumes and calibrated application equipment.

RESISTANCE: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product must conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

Mix Preparation

To prepare the application mixture:

1. Fill the spray tank with a portion of the required amount of water and begin agitation.
2. Add the specified amount of NUPRID® 4F Insecticide packets and allow packets to fully dissolve.
3. Fill the tank with the remaining water needed. Maintain sufficient agitation during mixing and application.

This product can be used with other pesticides and/or fertilizer solutions; refer to the Tank Mix and Compatibility Notes below. When tank mixtures of this product and other pesticides are involved, prepare the tank mixture as specified above and follow the suggested Mixing Order below.

This product may be applied by chemigation (see APPLICATION THROUGH IRRIGATION SYSTEMS section below) if allowed in the specific application sections.

TANK MIXES

Unless otherwise prohibited on this label or the label of an intended tank mix product, this product may be applied in combination with any pesticide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. **IMPORTANT: PESTICIDE TANK MIXES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL OR 24(c) LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.**

Compatibility

Before full-scale mixing of this product with other pesticides, fertilizers, secondary plant nutrients, adjuvants, surfactants or oils, you must determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent. **IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S**

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE () AMENTALS

SUPPLEMENTAL OR 24(c) LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

Mixing Order

When pesticide mixtures are needed, add wettable powders first, this product or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added and do not add an additional component until the previous is thoroughly mixed. A fertilizer/pesticide compatibility agent may be needed if a fertilizer solution is to be added to the mixture. Be sure to maintain constant agitation during both mixing and application to ensure uniformity of spray mixture. Further information on Tank Mixes is available from your local Nufarm representative.

APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION)

This product may be applied at rates specified on this label either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:100 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- Apply this product only through micro-irrigation (individual spaghetti tube), drip irrigation, overhead irrigation, and ebb and flood or hand-held or motorized calibrated irrigation equipment and only as directed in the specific directions. **DO NOT** apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- Be sure to remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system prior to application.
- A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES

If the source of water for your irrigation system is a public water supply, follow the instructions below:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.
- 8.

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE ORNAMENTALS

SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY

If the source of water for your irrigation system is NOT a public water supply, follow the instructions below:

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of material that is compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Water Volume

Chemigation applications of this product must be made as concentrated as possible. Retention of this product on target site of insect infestation is necessary for optimum activity. Chemigation of this product in water volumes exceeding 0.10 inch/Acre is not recommended.

Uniform Water Distribution and System Calibration

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift

DO NOT apply when wind speed favors drift beyond the area intended for treatment.

RESTRICTIONS

DO NOT graze treated areas or use clippings from treated areas for feed or forage.

Prevent runoff or puddling of irrigation water following application.

Keep children and pets off treated area until dry.

DO NOT apply this insecticide to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.

DO NOT apply more than 0.8 pts (12.8 fl oz) (0.4 lbs ai) per acre per year to outdoor plants/ornamentals.

Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

APPLICATION INSTRUCTIONS

NURSERY and GREENHOUSE GROWN ORNAMENTALS and VEGETABLE PLANTS

This product is for foliar and systemic insect control in and around field-grown nursery and container stock, indoor and outdoor ornamentals (including both greenhouse and interior landscapes) and ornamentals grown in flats, on benches or in beds. Apply this product by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests. Soil applications will result in translocation of the active ingredient upward into the plant system from root uptake. To assure optimum root uptake, apply product where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution has been shown to enhance the uptake of the active

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE ORNAMENTALS

ingredient. When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, soil applications must be made prior to anticipated pest infestation to achieve optimum levels of control.

Trees (including non-bearing fruit and nut), Shrubs, Evergreens, Flowers, Foliage Plants, Groundcovers, Interior Plantscapes, and Vegetable plants intended for resale only ¹		
FOLIAR APPLICATION		
Pests	Fluid ounces/ 100 gallons of water	
For control of: Larvae of: Adelgids Aphids Japanese beetles (adults) Lacebugs Leaf beetles (including Elm, Viburnum) Leafhoppers/Sharpshooters Leaf miners Mealybugs Sawfly larvae Whiteflies	0.75 (22 ml)	
For suppression of: Thrips		
Application Methods		
Mix product with the required amount of water and apply as desired dependent upon the selected use pattern. When making foliar applications on hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader/sticker is recommended. If concentrate or mist type spray equipment is used, apply an amount of product on the treatment area equivalent to the amount that would be used in a dilute application. This insecticide has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. The physical compatibility of this product may vary with different sources of pesticide products and local cultural practices. Any tank mixture which has not been previously tested must be prepared on a small scale (pint or quart jar), using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.		
Remarks		
Start treatments prior to establishment of high pest populations and reapply on an as needed basis.		
Restrictions		
¹ Only for use on vegetable plants intended for resale including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussel Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.		
BROADCAST APPLICATION		
Pests	Fluid ounces/1,000 FT ²	Fluid ounces/Acre
For control of: White grub larvae such as: Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle	0.23 – 0.30 (7.0 – 9.0 ml)	10.0 – 12.8 (0.625 – 0.8 pints)
Application Methods		
Mix required amount of product in sufficient water to uniformly and accurately cover the treatment area. DO NOT use less than 2 gallons of water per 1,000 sq ft		
Remarks		
For control of soil inhabiting pests, irrigate thoroughly to incorporate this insecticide into the upper soil profile. Bark Media: Media with 30% or more bark content may confer a shorter period of protection when treated with this product.		
Restrictions		
DO NOT apply more than 0.8 pints (12.8 fluid ounces) (0.4 lbs AI) per acre per year ¹ Only for use on vegetable plants intended for resale including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussel Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets Tomatillo, and Tomato.		

Trees, Shrubs, Flowers, and Groundcover	
SOIL APPLICATION (injection & drench)	
Pests	USE RATES
For control of: Adelgids Aphids Black vine weevil larvae Emerald ash borer ¹ Eucalyptus longhorned borer ¹ Flatheaded borers (including Bronze birch and Alder) ¹ Japanese beetles Lace bugs Leaf beetles (including Elm and Viburnum) Leafhoppers/Sharpshooters Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales White grub larvae Whiteflies	TREES: per inch of trunk diameter (DBH) 0.05 – 0.20 fluid ounces (1.5 – 6.0 ml) ²
	SHRUBS: per foot of shrub height 0.05 – 0.10 fluid ounces (1.5 – 3.0 ml)
	FLOWERS and GROUDCOVER: 0.23 - 0.30 fluid ounces (7.0 - 9.0 ml) / 1000 FT ²
For suppression of: Armored scales Thrips	Use the high rate
Application methods for TREES and SHRUBS	
Soil injection: Mix the required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per tree or shrub.	
Specific Soil Injection methods for trees and large shrubs:	
GRID System:	Holes must be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree.
CIRCLE System:	Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.
BASAL System:	Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.
Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree or shrub, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.	
Remarks	
¹ Application to trees already heavily infested with listed borers may not prevent the eventual loss of the trees due to existing pest damage and tree stress. ² Use higher specified rate for larger trees (over 8" D.B.H.) or for difficult to control insects or for trees with severe infestations.	
Restrictions	
DO NOT apply using Soil Injection methods in Nassau or Suffolk Counties of New York. DO NOT apply more than 0.8 pints (0.4 lbs AI) per acre per year.	
Application methods for FLOWERS and GROUNDCOVER	
Apply as a broadcast treatment and incorporate into the soil before planting or apply prior to bloom or after all flower petals have fallen off for established plants. If application is made to established plants, irrigate thoroughly after application. Bark Media: Media with 30% or more bark content may confer a shorter period of protection when treated with this product.	

GRASSY AREAS IN NURSERIES

Apply this product to grassy areas under and around field and container grown plants, on roadways and other grassy areas in and around nurseries to control soil inhabiting pests. Use this product for the suppression of Mole crickets and Cutworms. The active ingredient in this product has sufficient residual activity so that applications can be made prior to the egg laying activity of the target pest. Base the need for an application on historical monitoring of the site, previous records and experience, current season adult trapping, and other methods. Make applications prior to egg hatch of the target pests. Sufficient irrigation or rainfall is needed to facilitate the movement of active ingredient through the thatch.

Pests	Fluid ounces/1,000 FT ²	Fluid ounces/Acre
For control of: Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Chafers (Northern masked, Southern masked, European) Green June beetle Japanese beetle May or June beetle Oriental beetle <i>Phyllophaga spp.</i>	0.23 – 0.30 (7.0 – 9.0 ml)	10.0 – 12.8 (0.625 – 0.8 pints)
For control of: Mole crickets ¹	0.30 (9.0 ml)	12.8 (0.8 pints)
For suppression of: Chinchbugs ²		
Application Methods		
Apply this product in sufficient water to provide adequate distribution over the treatment area. The use of accurately calibrated equipment normally used for the application of soil insecticides is required. Use equipment which will produce a uniform, coarse spray droplet, use a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly. Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. DO NOT mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.		
Remarks		
For control of grubs, billbugs and annual bluegrass-weevil, make application prior to egg hatch of the target pest. Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific information regarding timing of application. ¹ For control of Mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, make a curative insecticide application with the application of this product. See tank mixing section above for general instructions on tank mixtures. ² For suppression of Chinchbugs, make application prior to or during the hatching of the first instar nymphs.		
Restrictions		
Maximum application rate per acre per year: 0.8 pints (12.8 fluid ounces) (0.4 lb of active ingredient) DO NOT make application when treatment area is waterlogged or soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treatment area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. DO NOT allow this product to contact plants in bloom if bees are foraging the treatment area.		

TURFGRASS (including Sod farms)

Use this product for the control of soil inhabiting pests of turfgrass. Use this product for suppression of cutworms and chinch bugs. Use as directed on turfgrass on sites such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms. The active ingredient in this product has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. Base the need for an application on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

SUB-LABEL B: NURSER\, GREENHOUSE and LANDSCAPE FUNDAMENTALS

Pests	Fluid ounces/1,000 FT ²	Fluid ounces/Acre
For control of: <u>Larvae of:</u> Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass atenius Chafers (Northern masked, Southern masked, European) European crane fly Green June beetle Japanese beetle May or June beetle Oriental beetle <i>Phyllophaga spp.</i>	0.23 – 0.30 (7.0– 9.0 ml)	10.0 – 12.8 (0.625 – 0.8 pints)
For control of: Mole crickets ¹	0.30 (9.0 ml)	12.8 (0.8 pints)
For suppression of: Chinchbugs ²		
Application Methods		
Apply this product in sufficient water to provide adequate distribution over the treatment area. The use of accurately calibrated equipment normally used for the application of soil insecticides is required. Use equipment which will produce a uniform, coarse spray droplet, use a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly. Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. DO NOT mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.		
Remarks		
For control of grubs, European crane fly, billbugs and annual bluegrass-weevil, make application prior to egg hatch of the target pest. Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific information regarding timing of application. ¹ For control of Mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, make a curative insecticide application with the application of this product. See tank mixing section above for general instructions on tank mixtures. ² For suppression of Chinchbugs, make application prior to or during the hatching of the first instar nymphs.		
Restrictions		
Maximum application rate per acre per year: 12.8 fluid oz (0.8 pints) (0.4 lb of active ingredient) DO NOT make application when treatment area is waterlogged or soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treatment area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. DO NOT allow this product to contact plants in bloom while bees are foraging the treatment area. DO NOT graze treated areas or use clippings from treated areas for feed or forage. DO NOT allow runoff or puddling of irrigation water following application. Keep children and pets off treated area until dry. Do not use for seed production.		

EBB & FLOOD APPLICATION

Apply this product through Ebb and Flood applications. To assure accurate uptake prior to treatment, bring a minimum of 10 plants up to a known field capacity and allow to dry out for one or two days. Re-wet these plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This will minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

Pests	Ornamental and vegetable plants ⁵ grown in containers		
	Pot size (inches)	Herbaceous species including vegetable plants ⁵ (1 or 2 plants/pot)	Woody perennials, Herbaceous species including vegetable plants ⁵ (3 or more/pot)
		ml / 100 plants	
For control of:			
Adelgids	2	0.80	1.25
Aphids			
Armored scales (suppression)	3	1.25	1.85
Fungus gnats (larvae only) ¹			
Japanese beetles (adults)	4	1.65	2.50
Lacebugs			
Leaf Beetles (including Elm and Viburnum)	5	2.10	3.15
Leafhoppers/Sharpshooters			
Leafminers	6	2.50	3.85
Mealybugs			
Psyllids	7	2.95	4.55
Root mealybugs ²			
Root weevil complex: (such as Apoka, Black vine, Citrus root) ³	8	3.30	5.00
Soft scales			
Thrips (suppression) ⁴	9	3.70	5.55
Whiteflies			
White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	10	4.15	6.25
	11	4.50	7.15
	12	5.00	8.35

¹ **Fungus gnat larvae** in the soil will be controlled by drench or incorporation. **No adult Fungus Gnat control.** Other foliar insect control is achieved by the uptake of this product from a healthy root system translocating the active ingredient up into the plant.

² **Root Mealybug** control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 1.0 fluid ounces (30 ml) in 150 gallons of water.

³ **Citrus Root Weevil:** For use on non-bearing citrus nursery stock.

⁴ **Thrips** suppression on foliage only. Thrips in buds and flowers will not be suppressed.

⁵ **Note:** For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape

DRENCH and IRRIGATION APPLICATIONS

Use this product for drench and irrigation application only on greenhouse and nursery grown ornamentals, vegetable plants intended for resale only, and interior plantscapes using soil drenches, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held or calibrated motorized irrigation equipment. This product may be applied at rates recommended on the label either alone or in tank mixtures with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:100 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation is necessary if the mixture is allowed to

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE FUNDAMENTALS

stand more than 24 hours. Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system. Only use this product through micro irrigation (individual spaghetti tubes), drip irrigation, overhead irrigation, ebb and flood, or handheld or motorized calibrated irrigation equipment. **DO NOT** apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non uniform distribution of treated water.

Pests	Ornamental and vegetable plants ⁵ grown in containers		
	Container size (inches)	Herbaceous species including vegetable plants ⁵ (1 or 2 plants/pot)	Woody perennials, Herbaceous species including vegetable plants ⁵ (3 or more/pot)
		# of Containers treated with 1.0 fluid oz (30 ml)	
For control of: Adelgids Aphids Fungus gnats (larvae only) ¹ Japanese beetles (adults) Lacebugs Leaf Beetles (including Elm and Viburnum leaf beetles) Leafhoppers (including glassy- winged sharpshooter) Leafminers Mealybugs Psyllids Root mealybugs ² Root weevil complex: (such as Apopka, Black vine, Citrus root weevils) ³ Soft scales Thrips (suppression) ⁴ Whiteflies White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	2	3000	2000
	3	2000	1350
	4	1500	1000
	5	1200	800
	6	1000	650
	7	850	550
	8	750	500
	9	675	450
	10	600	400
	11	550	350
	12	500	300
	Application method		
	Use sufficient volume to wet most of the potting medium without loss of liquid from the bottom of the container. Apply according to label directions. Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient due to leaching.		
	Ornamental and vegetable plants⁵ grown in flats, benches, or beds		
	0.34 fl. oz. (10 mL) per 1,000 square feet		
	Application method		
	Mix required amount in sufficient water to uniformly cover the area being treated. Do not use less than 2 gallons of mixture per 1,000 square feet. Apply as a broadcast treatment and incorporate into the medium before planting. Allow no leaching or runoff for 10 days after application.		
Remarks			

¹ **Fungus gnat larvae** in the soil will be controlled by drench or incorporation. **No adult Fungus Gnat control.** Other foliar insect control is achieved by the uptake of this product from a healthy root system translocating the active ingredient up into the plant.

² **Root Mealybug** control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 1.0 fluid ounces (30 ml) in 150 gallons of water.

³ **Citrus Root Weevil:** For use on non-bearing citrus nursery stock.

⁴ **Thrips** suppression on foliage only. Thrips in buds and flowers will not be suppressed.

⁵ **Note:** For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape

Pests	Containerized plants
-------	----------------------

SUB-LABEL B: NURSERY, GREENHOUSE and LANDSCAPE PLANTING

	Container Size (gallons)	# of Containers treated with 1.0 fluid oz (30 ml)
For control of: Adelgids Aphids Fungus gnats (larvae only) ¹ Japanese beetles (adults) Lacebugs Leaf Beetles (including Elm and Viburnum) Leafhoppers/Sharpshooters Leafminers Mealybugs Psyllids Root mealybugs ² Root weevil complex: (such as Apoka, Black vine, Citrus root) ³ Soft scales Thrips (suppression) ⁴ Whiteflies White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	1	244 – 340
	2	210 – 280
	3	185 – 220
	5	110 – 160
	7	75 – 100
	10	45 – 60
	15	30 – 40
	20	15 – 20
Application method		
Apply in sufficient water to wet the potting medium. Make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.		
Remarks		
¹ Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of this product from a healthy root system translocating the active ingredient up into the plant. ² Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 1.0 fluid ounces (30 mL) in 150 gallons of water. ³ Citrus Root Weevil: For use on non-bearing citrus nursery stock. ⁴ Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.		

Field and Forest Nurseries		
Pests	Fluid ounces / 1,000 ft of row	Fluid ounces / 1,000 square ft
For control of: White grub larvae ¹ (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	1.0 (30 ml)	0.34 (10 ml)
		12.8 Fluid ounces / Acre
Application method		
Apply as a uniform band on either side of the row using a band width six (6) inches wider than the actual root ball diameter to be dug. DO NOT allow bands in adjacent rows to overlap.		
Remarks		
Mowing of the vegetation in the area to be treated to a height of 3 inches or less prior to application will improve the consistency of control. Apply May through July. For target pest control, treatment must be followed by rainfall or irrigation. ¹ For grub control in areas of turf, apply as a broadcast application using 0.25 – 0.34 fluid ounces (7 – 10 ml) per 1,000 square feet (10.9 – 12.8 fluid ounces / Acre)		
Restrictions		
DO NOT use less than 2 gallons of spray volume per 1,000 square feet (85 GPA) DO NOT exceed 12.8 fluid ounces / acre per year (0.4 lbs AI/A)		

LANDSCAPE ORNAMENTALS and PLANTINGS

SUB-LABEL B: NURSER\ REENHOUSE and LANDSCAPE JAMENTALS

This product is for use on ornamentals and plantings in commercial and residential landscapes and interior plantscapes. It is a systemic product. Apply this product by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests. Soil applications will result in translocation of the active ingredient upward into the plant system from root uptake. Apply this product where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution has been shown to enhance the uptake of the active ingredient. When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, soil applications must be made prior to anticipated pest infestation to achieve optimum levels of control.

Trees, Shrubs, Evergreens, Flowers, Foliage Plants, Groundcovers, Interior Plantscapes		
FOLIAR APPLICATION		
Pests	Fluid ounces/ 100 gallons of water	
For control of: Larvae of: Adelgids Aphids Japanese beetles (adults) Lacebugs Leaf beetles (including Elm, Viburnum) Leafhoppers/Sharpshooters Leaf miners Mealybugs Sawfly larvae Whiteflies	0.75 (22 ml)	
For suppression of: Thrips		
Nuisance Ant Management: Use this product to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for nuisance ant populations. Applications can then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce ant populations.		
Application Methods		
Mix product with the required amount of water and apply as desired dependent upon the selected use pattern. When making foliar applications on hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader/sticker is recommended. If concentrate or mist type spray equipment is used, use an equivalent amount of product on the area sprayed, as would be used in a dilute application. This insecticide has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. The physical compatibility of this product may vary with different sources of pesticide products and local cultural practices. Any tank mixture which has not been previously tested must be prepared on a small scale (pint or quart jar), using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.		
Remarks		
Start treatments prior to establishment of high pest populations and reapply on an as needed basis.		
BROADCAST APPLICATION		
Pests	Fluid ounces/1,000 FT ²	Fluid ounces/Acre
For control of: White grub larvae such as: Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle	0.23 – 0.30 (7.0 – 9.0 ml)	10.0 – 12.8 (0.625 – 0.8 pints)
Application Methods		
Mix required amount of product in sufficient water to uniformly and accurately cover the treatment area. DO NOT use less than 2 gallons of water per 1,000 sq ft		
Remarks		
Irrigate thoroughly to incorporate this insecticide into the upper soil profile.		
Restrictions		
DO NOT apply more than 0.8 pints (0.4 lbs AI) per acre per year by application to outdoor ornamentals. DO NOT use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial production fruit and nut trees. DO NOT apply to Landscape Ornamentals and Plantings through any irrigation system. Follow application restrictions for Non-Agricultural Use Sites found in the RESTRICTIONS section of this label to		

protect bees and other insect pollinators.
Keep children and pets off treated area until dry.

Trees, Shrubs, Flowers, and Groundcover In Commercial and Residential Landscapes	
SOIL APPLICATION	
Pests	USE RATES
For control of: Adelgids Aphids Black vine weevil larvae Emerald ash borer ¹ Eucalyptus longhorned borer ¹ Flatheaded borers (including Bronze birch and Alder) ¹ Japanese beetles Lace bugs Leaf beetles (including Elm and Viburnum) Leafhoppers/Sharpshooters Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales White grub larvae Whiteflies	TREES: per inch of trunk diameter (DBH) 0.05 – 0.20 fluid ounces (1.5 – 6.0 ml) ²
	SHRUBS: per foot of shrub height 0.05 – 0.10 fluid ounces (1.5 – 3.0 ml)
	FLOWERS and GROUND COVER: 0.23 - 0.30 fluid ounces (7.0 – 9.0 ml) / 1000 FT ²
For suppression of: Armored scales Thrips	Use the high rate
Application methods for TREES and SHRUBS	
Soil injection: Mix the required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per tree or shrub.	
Specific Soil Injection methods for trees and large shrubs:	
GRID System:	Holes must be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree.
CIRCLE System:	Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.
BASAL System:	Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.
Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree or shrub, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.	
Remarks	
¹ Application to trees already heavily infested with listed borers may not prevent the eventual loss of the trees due to existing pest damage and tree stress. ² Use higher specified rate for larger trees (over 8" D.B.H.) or for difficult to control insects or for trees with severe infestations.	
Restrictions	
DO NOT apply using Soil Injection methods in Nassau or Suffolk Counties of New York. DO NOT apply more than 0.8 pints (0.4 lbs AI) per acre per year.	
Application methods for FLOWERS and GROUND COVER	
Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established prior to bloom or after all petals have fallen off. Irrigate immediately following application to established plants.	

Pomefruits In Commercial and Residential Landscapes: Apple, Crabapple, Loquat, Mayhew, Pear, Pear (oriental), Quince

FOLIAR APPLICATION		
Pests	USE RATES	
For control of: Aphids (except Woolly apple aphid) ¹ Leafhoppers/Sharpshooters ² Leafminers ³ Mealybugs ⁴ San Jose scale ⁵	0.75 fluid ounces (22 ml) / 100 gallons of water	3.0 fluid ounces/ Acre (90 ml)
Application methods		
Apply the specified dosage as a foliar spray as needed not to exceed 12.0 fluid ounces (0.09 lbs AI)/acre/year only after petal-fall is complete.		
Remarks		
The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. ¹ For control of Rosy apple aphid, apply prior to leafrolling caused by the pest. ² For late season (preharvest) control of Leafhopper species, apply this product while most Leafhoppers are in the nymphal stage. ³ For first generation Leafminer control, make 1 st application as soon as petal-fall is complete. Greatest Leafminer control will result from the earliest possible application. For 2 nd and succeeding generations of Leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A 2 nd application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. This product will not control late state larvae. ⁴ For control of Mealybugs, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the Mealybugs. ⁵ For San Jose scale, time applications to the crawler stage. Treat each generation.		
Restrictions		
DO NOT apply more than 3.0 fluid ounces (0.09 lbs AI) per acre in a single application. DO NOT make more than 4 applications. Allow 10 or more days between applications. Allow at least 7 days between last application and harvest. DO NOT use on Pomefruits grown for commercial production. DO NOT use in California for control on Pears. Follow application restrictions for Non-Agricultural Use Sites found in the RESTRICTIONS section of this label to protect bees and other insect pollinators. Keep children and pets off treated area until dry		

Pecan trees In Commercial and Residential Landscapes		
FOLIAR APPLICATION		
Pests	USE RATES	
For control of: Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	0.75 fluid ounces (22 ml) / 100 gallons of water	3.0 fluid ounces/ Acre (90 ml)
Application methods		
Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14-day interval may be required to achieve control. Scout and retreat if needed. Thorough uniform coverage of foliage is necessary for optimal control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage.		
Remarks		
The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.		
Restrictions		
DO NOT apply more than 9.0 fluid ounces (0.28 lbs AI) per acre per year. DO NOT make more than 3 applications per year. Allow 10 or more days between applications. DO NOT use on Pecans grown for commercial production. DO NOT use on Pecans in California unless directed by state approved 24(c) labeling. Follow application restrictions for Non-Agricultural Use Sites found in the RESTRICTIONS section of this label to protect bees and other insect pollinators. Keep children and pets off treated area until dry Allow at least 7 days between last application and harvest.		

Grapes In Commercial and Residential Landscapes		
FOLIAR APPLICATION		
Pests	USE RATES	
For control of: Leafhoppers/Sharpshooters Mealybugs	0.75 fluid ounces (22 ml) / 100 gallons of water	1.5 fluid ounces/ Acre (45 ml)
Application methods		
Apply specified dosage of this product as a foliar spray using 200 gallons of water per acre.		
Restrictions		
DO NOT apply more than 3.0 fluid ounces (0.09 lbs AI) per acre per year. . Allow 14 or more days between applications. Application can be made up to and including the day of harvest. Follow application restrictions for Non-Agricultural Use Sites found in the RESTRICTIONS section of this label to protect bees and other insect pollinators. Keep children and pets off treated area until dry		

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL [HANDLING]:

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the

container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable containers larger than 5 gallons]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[Refillable containers for return to Nufarm]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

SUB-LABEL B: NURSEF GREENHOUSE and LANDSCAPE FUNGICIDES

NUPRID is a registered trademark of Nufarm Americas Inc.

All other trademarks that appear on this label which are not owned by Nufarm Americas Inc. or its subsidiaries are the property of their respective owners.

(RV070914)

Urbanski, Jennifer

From: carrie.tackema@us.nufarm.com
Sent: Wednesday, July 09, 2014 8:38 AM
To: Urbanski, Jennifer
Subject: RE: 228-528
Attachments: 000228-00528.20140709.draft.pdf

Here you go

Best regards,
Carrie



Nufarm
Carrie M. Tackema
Regulatory Manager
Nufarm Americas, Inc.
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560

☎: (919) 379-2528 (Office)
☎: (919) 323-1368 (Cell)
☎: (919) 467-5923 (Fax)
✉: carrie.tackema@us.nufarm.com

THIS MESSAGE IS INTENDED ONLY FOR THE USE OF THE INDIVIDUAL OR ENTITY TO WHICH IT IS ADDRESSED AND MAY CONTAIN INFORMATION THAT IS PRIVILEGED, CONFIDENTIAL AND EXEMPT FROM DISCLOSURE. If the reader of this message is not the intended recipient or an employee of agent responsible for delivering the message to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this communication is strictly prohibited. If you have received this communication in error, please delete, destroy and in doing so do not save the message. Nufarm Americas, Inc. and its affiliated companies. Fax 919 342 5176

From: "Urbanski, Jennifer" <urbanski.jennifer@epa.gov>
To: "carrie.tackema@us.nufarm.com" <carrie.tackema@us.nufarm.com>,
Date: 07/09/2014 06:24 AM
Subject: RE: 228-528

Hi Carrie, just a couple of things from the PM and we'll be all set!

- 1) On page 21-22, why is there a lighter shade of gray in the box? The text indicates that the shaded rates may not provide residual control (and I assume this means the dark gray cells), but it doesn't say anything about the light gray. Remove the light gray shading if it doesn't mean anything but if it does, explain it in the text below the table.
- 2) On page 31, change "DO NOT apply in Almonds" to "DO NOT apply on Almonds".
- 3) On page 52, delete "broadcast" from "... per acre per year by broadcast application to outdoor ornamentals".
- 4) On page 54, change "Apply the specified dosage as a foliar spray as needed after petal-fall is complete." to read "Apply the specified dosage as a foliar spray as needed (not to exceed [add max rate per year] only after petal-fall is complete."

From: carrie.tackema@us.nufarm.com [<mailto:carrie.tackema@us.nufarm.com>]

Sent: Tuesday, July 08, 2014 12:06 PM

To: Urbanski, Jennifer

Subject: Re: 228-528

Hi Jenn, I thought this might happen. I originally sent the poultry amendment to Julie Chao, because I was working with UPI to get the change made like they did to their label, and Julie worked on theirs. So yes, the two amendments have crossed over. I have updated the label to add the additional language on page 53.

Best regards,
Carrie



Carrie M. Tackema
Regulatory Manager
Nufarm Americas, Inc.
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560

☎: (919) 379-2528 (Office)

☎: (919) 323-1368 (Cell)

☎: (919) 467-5923 (Fax)

✉: carrie.tackema@us.nufarm.com


THIS MESSAGE IS INTENDED ONLY FOR THE USE OF THE INDIVIDUAL OR ENTITY TO WHICH IT IS ADDRESSED AND MAY CONTAIN INFORMATION THAT IS PRIVILEGED, CONFIDENTIAL AND EXEMPT FROM DISCLOSURE. If the reader of this message is not the intended recipient or an employed of agent responsible for delivering the message to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this communication is strictly prohibited. If you have received this communication in error, please delete, destroy and in doing so do not save the message. Nufarm Americas, Inc. and its affiliated companies. Fax 919 342 5176

From: "Urbanski, Jennifer" <urbanski.jennifer@epa.gov>
To: "carrie.tackema@us.nufarm.com" <carrie.tackema@us.nufarm.com>,
Date: 07/08/2014 11:23 AM
Subject: 228-528

Hey Carrie, just got this amendment to change the poultry house restriction. You also included the containerized citrus rates that have already been reviewed and approved. The label looks good, just one comment: On page 53, after "...before planting or apply after plants are established", add "prior to bloom or after all petals have fallen off". Thanks!

Jennifer Urbanski, Ph.D., Biologist
Insecticide-Rodenticide Branch, S7221
Registration Division (7505P)
U.S. Environmental Protection Agency
1200 Pennsylvania Ave. NW
Washington, DC 20460
(703) 347-0156

EXPEDITE

 EPA	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other:	OPP Identifier Number
---	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 228-528	2. EPA Product Manager Venus Eagle	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Nuprid 4F Insecticide	PM# 1	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 11901 S. Austin Avenue Alsip, IL 60803 <u>Please send all correspondence to "contact point" listed below</u>		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(1), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

- | | |
|--|--|
| <input type="checkbox"/> Amendment – Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - Explain below |

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)


Application for Label Amendment

CONTACT: **Carrie M. Tackema (919) 379-2528 or carrie.tackema@us.nufarm.com**
Nufarm Americas, Inc.
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt. No. per container	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other
* Certification must be submitted			
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container 16, 32, 64 fl. oz.; 1, 2.5, 220, 250 gal.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)

Name Carrie M. Tackema	Title Regulatory Manager	Telephone No. (Include Area Code) (919) 879-2528
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Manager	
4. Typed Name Carrie M. Tackema	4. Date June 16, 2014	



+1 919.379.2510
+1 919.467.5923
4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560
www.nufarm.com

Via Courier Delivery

June 16, 2014

Ms. Venus Eagle, PM #1
USEPA (7504P)
Document Processing Desk (AMEND)
Room S4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

**Subject: Nuprid® 4F Insecticide, EPA Reg. No. 228-528
Application for Label Amendment**

Dear Ms. Eagle:

Enclosed please find a label amendment for the above-referenced product. The purpose of this amendment is to revise the Poultry Housing section, specifically, removal of the 7-day restriction. This will support the registration and match the CPS label for EPA Reg. No. 34704-931. Further, this action incorporates the recent changes submitted under NOTIF action to revise the use rates in Citrus (submitted to your attention on May 27, 2014). Nufarm respectfully requests an EXPEDITED REVIEW of the proposed labeling. Also enclosed in support of this Application are the following:


- Application for Pesticide Registration - AMEND;
- Label certification; and
- Two (2) copies of proposed product labeling for each product (one copy with changes highlighted, and one in pdf format on CD-rom).

Should you have any questions or require additional information, please contact me at (919) 379-2528 or by email at carrie.tackema@us.nufarm.com.

Sincerely,

Carrie M. Tackema
Regulatory Manager

Enclosure(s)





Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803
Phone: 708-377-1330 Fax: 708-377-1333

4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560
Phone: 919-379-2510 Fax: 919.467.5923
www.us.nufarm.com

Via Courier Delivery

June 9, 2014

Ms. Venus Eagle, PM #1
USEPA (7504P)
Document Processing Desk
Room S4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

Subject: Final Printed Labeling
Nuprid® 4F MAX Insecticide, EPA Reg. No. 228-528

Dear Ms. Eagle:

Enclosed please find final printed labeling for the above-referenced product.


Should you have any questions or require additional information, please contact me at (919) 379-2528 or by email at carrie.tackema@us.nufarm.com.

Sincerely,

A handwritten signature in black ink, appearing to be 'Carrie M. Tackema', written over a horizontal line.

Carrie M. Tackema
Regulatory Manager

Enclosure(s)



EXPEDITE

EPA United States Environmental Protection Agency Washington, DC 20460		<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other:	OPP Identifier Number
Application for Pesticide - Section I			
1. Company/Product Number 228-528		2. EPA Product Manager Venus Eagle	
4. Company/Product (Name) Nuprid 4F Insecticide		3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 11901 S. Austin Avenue Alsip, IL 60803 <u>Please send all correspondence to "contact point" listed below</u>		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	
Section - II			
<input type="checkbox"/> Amendment – Explain below. <input checked="" type="checkbox"/> Final printed label <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Notification - Explain below. <input type="checkbox"/> Other - Explain below			
Explanation: Use additional page(s) if necessary. (For Section I and Section II.) <div style="text-align: center;">FINAL PRINTED LABELING</div> <p>CONTACT: <u>Carrie M. Tackema (919) 379-2528 or carrie.tackema@us.nufarm.com</u> Nufarm Americas, Inc. 4020 Aerial Center Parkway, Suite 101 Morrisville, NC 27560</p>			
1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt. No. per container	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other...
*Certification must be submitted			
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 16, 32, 64 fl. oz.; 1, 2.5, 220, 250 gal.	
		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			
Section - IV			
1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)			
Name Carrie M. Tackema		Title Regulatory Manager	
		Telephone No. (Include Area Code) (919) 379-2528	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title Regulatory Manager	
4. Typed Name Carrie M. Tackema		4. Date June 9, 2014	

NOT REVIEWED
In Accordance with PR Notice 82-2
Based on Draft Labeling Dated

GROUP 4A INSECTICIDE

Nuprid[®]

4F Max

INSECTICIDE

A systemic and foliar insecticide for use on listed field crops including cotton, tobacco, potato, soybeans and peanuts; in citrus, tree nut, and fruit orchards; on field and greenhouse vegetables; on berry, bush and vine crops; and on other listed crops

ACTIVE INGREDIENT:

Imidacloprid, 1 -[(6-Chloro-3-pyridinyl)methyl]-
N-nitro-2-imidazolidinimine

40.4%

OTHER INGREDIENTS:

59.6%

TOTAL:

100.0%

Contains 4 pounds of imidacloprid per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE INSIDE BOOKLET FOR FIRST AID AND
ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or
Exposure, Call CHEMTREC
(800) 424-9300

For Medical Emergencies Only,
Call (877) 325-1840

EPA Reg. No. 228-528

EPA Est. No. indicated by the first two letters of the batch number on this package
(AL) 228-IL-002, (YU) 67545-AZ-001

Net Contents

1 Gal.
(3.78 L)

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803



file: 12368-4_book_art
folder: 12368-4_Nufarm_12815000
colors: CMYK (processed to match 348, 654)
DO NOT ALTER PERCENTAGES FOR 654 BLUE

created: 05-24-10 jw
alt: 03-05-14 jw
size: 4.75"(w) x 4.75"(h)
☐ dp proofed

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid contact with eyes or clothing. Wear protective eye wear. Wear long sleeved shirt and long pants, socks, shoes, and chemical resistant gloves made of waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

FIRST AID	
IF INHALED	<ul style="list-style-type: none"> • Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN No specific antidote is available. Treat the patient symptomatically.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS


User must:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. **DO NOT** apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates. This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

(continued)

PROTECTION OF POLLINATORS *(continued)*

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.


Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product, by any application method, to linden, basswood or other *Tilia* species in the State of Oregon.

See individual crops for specific pollinator protection application restrictions. If none exist under the specific crop, for foliar applications, follow these directions for use for crops that are contracted to have pollinator services or for food/feed and commercially grown ornamentals that are attractive to pollinators:




1. FOR CROPS UNDER CONTRACTED POLLINATION SERVICES

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless the following condition has been met:

If an application must be made when managed bees are at the treatment site, the beekeeper providing the pollination services must be notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.





2. FOR FOOD CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

PRODUCT INFORMATION

Thorough uniform coverage is necessary to achieve insect control. A spray adjuvant may be used to improve coverage. This product may not knockdown established and heavy insect populations with a single application. Two applications may be required to achieve control; retreat if needed and as directed on this label. This product may be tank mixed with other insecticides as specified for knockdown of pests or for improved control of other pests.

Applying this product to crops grown for production of true seed intended for private or commercial planting is not permitted unless allowed under state approved 24(c) labeling. Additional information on this product uses for listed crops and other questions may be obtained from the Cooperative Extension Service, PCA's, consultants or your local Nufarm representative.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product must conform to resistance management strategies established for your use area.

This product contains a Group 4A insecticide called imidacloprid. Insect biotypes with acquired or inherent tolerance to group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control of targeted species. This may eventually result in partial or total loss of control of those species by this product and other Group 4A products.

The active ingredient in this product is a member of neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of this product and/or other Group 4A products having the same or similar mode of action.

Following a neonicotinoid block of treatments, Nufarm strongly encourages the rotation to a block of applications with effective products of a different mode of action before using additional applications of neonicotinoid products. Use of a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect's pest's ability to develop resistance to this class of chemistry.

Foliar applications of this product or other Group 4A products from the neonicotinoid chemical class must not be used on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Intruder, Impulse, Leverage, Pasada, Provado, Trimax Pro and Venom. Other 4A Group, neonicotinoid products used as soil/seed treatment include: Admire Pro, Advise, Alias, Belay, Clutch, Couraze, Cruiser, Gaucho, Macho, Macho Max, Platinum, Venom and Widow. Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

PRODUCT USE INSTRUCTIONS

OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

SPRAY DRIFT MANAGEMENT

The responsibility of avoiding spray drift is with the applicator. The applicator must consider weather related factors and the interaction of application equipment when making application decisions.

Mixing and Loading Requirements

The use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well head, sinkholes or field drains.

Aerial Applications

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size

The droplet size is an important factor and can influence drift. Small droplets (<150 – 200 microns) drift to a greater extent than large droplets. Applications typically should be made to deliver the largest droplet range that provides adequate control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. **DO NOT** apply when winds are greater than 15 mph and avoid gusty and windless conditions.

Restrictions During Temperature Inversions

DO NOT make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions typically restrict vertical air mixing, which then could cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions typically are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. The following specific drift management practices must be followed.

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- **DO NOT** allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

No-Spray Zone Requirements for Foliar Applications

DO NOT apply by ground within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

RUNOFF MANAGEMENT

DO NOT cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff must be employed.

ENDANGERED SPECIES NOTICE

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

ROTATIONAL CROPS

Crops which are listed on imidacloprid labels or crops that have existing tolerances for imidacloprid may be planted in treated areas as soon as practical after the last imidacloprid application. Crops that are not found on an imidacloprid label, or crops that do not have existing tolerances for imidacloprid, may not be planted in treated areas for 12-months after the last application. Refer to the table below for plantback intervals for different crops. Note that if cover crops are planted any time after an application of this product, those crops may not be grazed or harvested for food or feed.

ROTATIONAL CROPS - PLANT-BACK INTERVALS	
IMMEDIATE PLANT-BACK:	All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop & sweet), rapeseed, sorghum, sugar beet and wheat.
30-DAY PLANT-BACK:	Cereals (including buckwheat, millet, oats, rice, rye, and triticale), soybeans and safflower
10-MONTH PLANT-BACK:	Onions and bulb vegetables
12-MONTH PLANT-BACK:	All other crops

APPLICATION INSTRUCTIONS

This product can be applied as a foliar spray, or as a soil treatment (see Crop Specific Restrictions and Limitations). Thorough uniform coverage is necessary to achieve insect control. Use adequate spray volumes, properly calibrated application equipment, and an adjuvant to improve coverage. Failure to provide adequate coverage and retention of this product on leaves and fruit may result in loss of insect control or delay in onset of activity.

This product may not knockdown established and heavy insect populations with a single application. Two applications may be required to achieve control. Scout fields and retreat if needed.

This product may be tank mixed with other insecticides as recommended for knockdown of pests or for improved control of other pests.

Apply this product with properly calibrated ground or aerial application equipment.

Minimum spray volumes unless otherwise specified on the Crop Specific Restrictions and Limitations section are 10 gallons/Acre by ground applications and 5 gallons/Acre through aerial equipment.

This product may also be applied by overhead chemigation (see **CHEMIGATION APPLICATION** section below) if allowed in crop specific recommended application section.

Apply specified rate per acre as a foliar spray as pest populations begin to build. **DO NOT** apply more than 0.5 lbs. active ingredient per acre, per year, regardless of formulation or method of application, unless specified within the **Crop Specific Restrictions and Limitations** for a given crop.

Mix Preparation

To prepare the application mixture:

1. Fill the spray tank with a portion of the required amount of water and begin agitation.
2. Add the specified amount of NUPRID® 4F Insecticide.
3. Fill the tank with the remaining water needed. Maintain sufficient agitation during mixing and application.

NOTE: This product may also be used with other pesticides and/or fertilizer solutions; refer to the Tank Mix and Compatibility Note below. When tank mixtures of this product and other pesticides are involved, prepare the tank mixture as specified above and follow the suggested Mixing Order below.

Tank Mixes

Unless otherwise prohibited on this label or the label of an intended tank mix product, this product may be applied in combination with any pesticide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. **IMPORTANT: PESTICIDE TANK MIXES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL OR 24(C) LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.**

Compatibility

Before full-scale mixing of this product with other pesticides, fertilizers, secondary plant nutrients, adjuvants, surfactants or oils, you must determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent. **IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL OR 24(c) LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.**

Mixing Order

When pesticide mixtures are needed, add wettable powders first, this product or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added and do not add an additional component until the previous is thoroughly mixed. A fertilizer/pesticide compatibility agent may be needed if a fertilizer solution is to be added to the mixture. Be sure to maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

Further information on Tank Mixes is available from your local Nufarm representative.

CHEMIGATION APPLICATION

Types of Irrigation Systems

Chemigation applications of this product may be made to crops through overhead sprinkler chemigation systems if specified in crop-specific recommendation sections. **DO NOT** apply this product through any other type of irrigation system.

Water Volume

Chemigation applications of this product must be made as concentrated as possible. Retention of this product on target site of insect infestation is necessary for optimum activity. Chemigation of this product in water volumes exceeding 0.10 inches/Acre are not recommended.

Uniform Water Distribution and System Calibration

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift

DO NOT apply when the wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or normally shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

CROP-SPECIFIC RESTRICTIONS AND LIMITATIONS

FIELD CROPS

COTTON		
SOIL APPLICATIONS		
Pests	Fluid ounces/1,000 row-feet	Fluid ounces/Acre
For control of: Cotton aphid Plant bugs Thrips Whiteflies	0.65	8.5 – 10.6 (Depending on row-spacing)
Application Methods		
Apply specified dosage by one of the following methods: 1. In-furrow or narrow band spray. When applying as an in-furrow spray, direct application on or below the seed at planting; OR 2. Narrow band application below the eventual seed bed row in a bedding operation 7 or fewer days before planting; OR 3. Chemigation into root zone through low-pressure drip or trickle irrigation equipment.		
Restrictions		
Maximum amount of product allowed per year: 10.6 fluid ounces/Acre (0.33 lb AI/Acre) DO NOT apply more than 0.5 lbs. of active ingredient per acre per year of NUPRID® 4F Insecticide, Provado®, Trimax® or Leverage®, including seed treatment as Gaucho®, soil and foliar uses. DO NOT graze treated fields after any application of this product. See Resistance Management section of this label.		
FOLIAR APPLICATIONS		
Pests	Fluid ounces/Acre	
For control of: Cotton aphid Cotton fleahopper Bandedwinged whitefly Plant bugs (excludes <i>Lygus hesperus</i>) Green stink bug Southern green stink bug Bollworm/Budworm (ovicidal effect)	1.0 – 2.0	
For suppression of: Lygus bugs (<i>Lygus hesperus</i>) Whiteflies (other than banded winged whitefly)	2.0	

(continued)

COTTON (continued)**Application Methods**

Apply through properly calibrated ground, aerial, or chemigation application equipment.

Restrictions

Pre-Harvest Interval (PHI): **14 days**

Minimum interval between applications: **7 days**

Maximum amount allowed per year: **10.0 fluid ounces/Acre** (0.31 lb. AI/A)

DO NOT apply more than 0.5 lbs. of active ingredient per acre per year of NUPRID® 4F Insecticide, Provado®, Trimax® or Leverage®, including seed treatment as Gaucho®, soil and foliar uses.

DO NOT graze treated fields after any application of this product.

TANK MIX APPLICATIONS

Pests (in addition to those listed above)	NUPRID 4F Rate Fluid ounces/Acre	BIDRIN 8 Rate¹ Fluid ounces/Acre
For early season control of: Thrips	1.0 – 1.5	1.6 – 3.2
For mid to late season control of: Plant bugs Stink bugs (including Brown stink bug) Grasshoppers Saltmarsh caterpillar Cotton leafperforator		4.0 – 8.0

Remarks

This product can be tank mixed with other pesticides and/or fertilizer solutions. When tank-mixing this product with other pesticides, prepare the tank mixture as specified above in the **Mix Preparation** instructions section. Follow the following general mixing order:

1. Add wettable powders first;
2. Add this product or other flowables second;
3. Add emulsifiable concentrates last.

Be sure to maintain agitation as each component is added and do not add an additional component until the previous component is thoroughly integrated into the mixture. If a fertilizer solution is added, a fertilizer-pesticide compatibility agent may be needed.

Restrictions (in addition to those listed above for NUPRID 4F foliar applications)

¹ Refer to the BIDRIN 8 product label for specific use instructions; observe all use restrictions and precautions that appear on the BIDRIN 8 label.

POTATO**SOIL APPLICATIONS**

Pests	Fluid ounces/1,000 row-feet	Fluid ounces/Acre
For control of: Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid For suppression of: Wireworms (with in-furrow spray at planting) For suppression of disease symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis (PLRV)	0.45 – 0.65	6.5 – 10.0

Application Methods

Apply specified dosage in one of the following methods:

1. In-furrow spray during planting directed on seed pieces or seed potatoes; OR
 2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil; OR
 3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil; OR
 4. Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting.
- For effective pest control or suppression, applications of this product must be placed below soil surface and in contact with seed piece or within root zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of this product may be made in a 2 to 4 inch band (width of planter shoe opening) and completely covered.

Restrictions

Maximum amount of product allowed per year: **10.0 fluid ounces/Acre** (0.31 lb AI/Acre)

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Colorado Potato beetle Flea beetles Fleahoppers Psyllids	1.5

(continued)

POTATO (continued)**Application Methods**

Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum amount allowed per year: **6.4 fluid ounces/Acre** (0.20 lb. AI/A)

SEED PIECE APPLICATIONS¹

Pests	Fluid ounces/100 lbs seed	Fluid ounces/Acre
For control of: Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid Wireworms (seed piece protection)	0.2 – 0.4	4.0 – 8.0
For suppression of disease symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis (PLRV)	0.4	8.0

Application Methods

Apply specified dosage as a diluted spray onto seed pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part this product. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after this product's application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed pieces as soon as possible after treating avoiding prolonged exposure of seed pieces treated with this product to sunlight and in accordance with the recommendation of your local Extension specialist. Consult your local Nufarm representative or crop protection product dealer for information relevant to your area.

Remarks

¹ Based on a seeding rate of 2000 lbs/Acre

Restrictions

Maximum amount of product allowed per year: **10.0 fluid ounces/Acre** (0.31 lb AI/Acre)

DO NOT use treated seed pieces for food, feed, or fodder.

DO NOT apply any subsequent applications of this product (in-furrow), Gaucho®, Leverage® or Provado® following a seed-piece treatment of this product.

Apply only in areas that are equipped to remove spray mist or dust or with adequate ventilation.

SOYBEAN**FOLIAR APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: Aphids Bean leaf beetle Cucumber beetles / Rootworm adults Japanese beetle (adults) Leafhoppers Whiteflies	1.5
Application Methods	
Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 21 days Minimum interval between applications: 7 days Maximum amount allowed per year: 3.65 fluid ounces/Acre (0.13 lb. AI/A) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	

TOBACCO**SOIL APPLICATIONS**

Pests	Fluid ounces/1,000 plants (as seedling tray drench)	Fluid ounces/1,000 plants (in-furrow or transplant-water)
For control of: Aphids Flea beetles	0.5	0.7
For control of: Mole crickets Whiteflies Wireworms	0.7 – 1.4	0.9 – 1.4
For suppression of disease symptoms of: Tomato spotted wilt virus (TSWV)	1.4	1.4

(continued)

TOBACCO *(continued)***Application Methods**

Apply specified dosage of this product in one of the following methods:

1. Uniform, broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting, followed immediately by overhead irrigation to wash this product from foliage into potting media. Failure to wash this product from foliage may result in a reduction in pest control. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.; OR
2. In-furrow spray or transplant-water drench during setting.; OR
3. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Remarks

Important Note: Proper tray drench applications of this product have been shown to be the most efficacious method of application. However, the specified rate of this product may be applied as a combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of this product into the plant and a delay in control.

Restrictions

Pre-Harvest Interval (PHI): **14 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids	0.8 – 1.6
For control of: Flea beetles Japanese beetles	1.6

Application Methods

Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **14 days**

Minimum interval between applications: **7 days**

Maximum amount allowed per year: **9.0 fluid ounces/Acre** (0.28 lb. AI/A)

PEANUT	
SOIL APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	8.0 – 12.0
For suppression of: Thrips	12.0
Application Methods	
Apply as a: 1. Chemigation into root-zone through properly calibrated low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.; OR 2. In-furrow spray directed on or below seed.	
Remarks	
Applications of this product have been shown to increase the incidence of Tomato spotted wilt virus (TSWV), and possibly other tospoviruses, on multiple varieties of peanut. Prior to making product applications, contact the State, Cooperative This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling. Extension Service, or Nufarm representative, for recommendations to discuss the risk and benefits of imidacloprid applications.	
Restrictions	
Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	1.4
Application Methods	
Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 14 days Minimum interval between applications: 5 days Maximum amount allowed per year: 4.2 fluid ounces/Acre (0.13 lb. AI/A) ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	

VEGETABLE CROPS

CUCURBIT VEGETABLES: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*).

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Cucumber beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	8.0 – 12.0
For suppression of disease symptoms of: Bacterial wilt (as vectored by various cucumber beetles) Leaf silvering resulting from whitefly feeding	12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray directed on or below seed; OR
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR
5. Post-seeding drench, transplant-water drench, or hill drench; OR
6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

(continued)

CUCURBIT VEGETABLES: *(continued)***GREENHOUSE APPLICATIONS¹**

Pests	Fluid ounces/1000 plants
For control of: Aphids Whiteflies	0.05

Application Methods

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash this product from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash this product from foliage may result in reduced pest control; OR
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

Remarks

The application made in the planthouse is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.

Important Note: Not all varieties of cucurbit vegetables have been tested for tolerance to this product applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

Restrictions

Maximum number of greenhouse applications: **1**

Maximum amount of product allowed per greenhouse application: **0.05 fluid ounce** (0.001568 lb AI)/**1,000 plants**.

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

FRUITING VEGETABLES: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo.

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Colorado potato beetles Flea beetles Leafhoppers Thrips (foliage-feeding only) Whiteflies	Okra & Pepper 8.0 – 16.0 Other Listed Crops 8.0 – 12.0
For suppression of disease symptoms of: Tomato mottle virus Tomato spotted wilt virus Tomato yellow leaf curl virus	

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray directed on or below seed; OR
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR
5. Post-seeding drench, transplant-water drench, or hill drench; OR
6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum amount of product allowed on Okra and Pepper per crop season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre).

Maximum amount of product allowed on other listed fruiting vegetable crops per application: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre).

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

FOLIAR APPLICATIONS (FIELD)

Pests	Fluid ounces/1000 plants
For control of: Aphids Colorado potato beetle Leaf beetles Whiteflies ¹	1.5 – 2.5

(continued)

FRUITING VEGETABLES: *(continued)*

Pests	Fluid ounces/1000 plants
For control of: Pepper weevil (Pepper only) ²	2.5

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Remarks

Applications of this product must be incorporated into a full-season program, where alternations of effective products from multiple classes of chemistry and different modes of actions are utilized in a blocked or windowed approach. For additional information, please contact your Nufarm representative, Extension Specialist or crop advisor.

¹Higher specified rate within the rate range must be used when targeting adult whiteflies.

²For pepper weevil, apply specified dosage of this product by ground equipment only. Time applications prior to a damaging pest population becoming established. Good coverage of foliage and fruit is necessary for target pest control.

Restrictions

Pre-Harvest Interval (PHI): **0 days**

Minimum interval between applications: **5 days**

Maximum amount of product allowed per crop season: **7.7 fluid ounce (0.24 lb AI)/A**

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

GREENHOUSE APPLICATIONS¹

Pests	Fluid ounces/1000 plants
For control of: Aphids Whiteflies	0.05

Application Methods

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash this product from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash this product from foliage may result in reduced pest control; OR
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

(continued)

FRUITING VEGETABLES: *(continued)***Remarks**

The application made in the planthouse is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.

Important Note: Not all varieties of cucurbit vegetables have been tested for tolerance to this product applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

Restrictions

Maximum number of greenhouse applications allowed: 1

Maximum amount of product allowed per greenhouse application: **0.05 fluid ounce** (0.001568 lb AI)/1,000 plants. **DO NOT** use on crops grown for seed unless allowed by state approved 24(c) labeling.

¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

GREENHOUSE VEGETABLES: Mature Cucumber and Tomato plants in production greenhouses ONLY.**Pests****Fluid ounces/1000 plants****For control of:**

Aphids
Whiteflies

0.7

Application Methods

Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. **DO NOT** apply to immature plants since phytotoxicity may occur.

Remarks

Make application when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (*Onius* sp.) can occur when this product is applied.

Many varieties of vegetables have been tested for tolerance to this product and show good safety. However, certain varieties may show more sensitivity to this product. Therefore, treat a few plants before treating the whole greenhouse.

Restrictions

Pre-Harvest Interval (PHI): **0 days**

Maximum number of applications per crop season: 1

Maximum amount of product allowed per crop season: **0.7 fluid ounce** (0.022 lb. AI)/1,000 plants.

DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.

GLOBE ARTICHOKE**SOIL APPLICATION¹**

Pests	Fluid ounces/1000 plants
For control of: Aphids Leafhoppers	8.0 – 16.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray directed on or below seed.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounce/Acre** (0.50 lb. AI/A)

¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATION

Pests	Fluid ounces/1000 plants
For control of: Aphids Leafhoppers	1.6 – 4.0

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum Interval between applications: **14 days**

Maximum amount of product allowed per year: **16.0 fluid ounce/Acre** (0.50 lb. AI/A)

HERBS: Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

SOIL APPLICATIONS¹ (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	8.0 – 12.0
For suppression of: Thrips (foliage-feeding only)	

Application Methods

Apply specified dosage in one of the following methods:

1. In-furrow spray during planting directed on or below seed; OR
2. In-furrow spray or transplant-water drench during setting or transplanting; OR
3. Shankd-into or below eventual seed-line; OR
4. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Remarks

Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat only small areas or numbers of plants and evaluate prior to full-scale use.

Restrictions

Pre-Harvest Interval (PHI): **14 days**

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Leaf beetles Leafhoppers Whiteflies	1.4

(continued)

HERBS: (continued)**Application Methods**

Apply as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Remarks

The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage and control.

Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, treat only small areas or numbers of plants and evaluate prior to full-scale use.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **5 days**

Maximum amount of product allowed per crop season: **4.2 fluid ounce** (0.13 lb AI/A)

HEAD and STEM BRASSICA VEGETABLES¹: Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gailon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, plus Turnip tops (leaves).

LEAFY GREENS VEGETABLES¹: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Raddicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), **Watercress²** (including upland).

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre (on 36 in. rows)
For control of: Aphids Leafhoppers Thrips (foliage-feeding only) Whiteflies	5.0 – 12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray directed on or below seed; OR

(continued)

LEAFY GREENS VEGETABLES¹: *(continued)***Application Methods** *(continued)*

3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR
5. Post-seeding drench, transplant-water drench, or hill drench; OR
6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.

RestrictionsPre-Harvest Interval (PHI): **21 days**Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)¹ **DO NOT** use on crops grown for seed unless permitted by state approved 24(c) labeling.² For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following application. Applications must be made to fully leafed-up canopies only. **DO NOT** apply to native cress growing in streams or other bodies of water.**FOLIAR APPLICATIONS (FIELD)**

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	1.5

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

RestrictionsPre-Harvest Interval (PHI): **7 days**Minimum interval between applications: **5 days**Maximum amount of product allowed per crop season: **7.7 fluid ounces/Acre** (0.23 lb. AI/A)¹ **DO NOT** use on crops grown for seed unless permitted by state approved 24(c) labeling.² For applications made to watercress, production fields must be drained of water at least 24 hours prior to application and water must not be reapplied to the field for a minimum of 24 hours following application. Applications must be made to fully leafed-up canopies only. **DO NOT** apply to native cress growing in streams or other bodies of water.

LEAFY PETIOLE VEGETABLES: Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, finocchio), Rhubarb, Swiss chard

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre (on 36 in. rows)
For control of: Aphids Leafhoppers Thrips (foliage-feeding only) Whiteflies	5.0 – 12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray directed on or below seed; OR
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours of application; OR
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; OR
5. Post-seeding drench, transplant-water drench, or hill drench; OR
6. Subsurface side-dress on both sides of each row. This product must be incorporated into root-zone.

Restrictions

Pre-Harvest Interval (PHI): **45 days**

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

LEGUME VEGETABLES (except Soybean, dry):

Edible podded and Succulent shelled pea and Bean and Dried Shelled Pea and Bean including:

Bean - *Lupinus* spp. (grain lupin, sweet lupin, white lupin, and white sweet lupin)

Bean - *Phaseolus* spp. (field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean) Bean - *Vigna* spp. (adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

Pea - *Pisum* spp. (dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

Other Beans and Peas - Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean.

SOIL APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	8.0 – 12.0

LEGUME VEGETABLES (except Soybean, dry): *(continued)***SOIL APPLICATIONS (FIELD)** *(continued)*

Pests	Fluid ounces/Acre
For suppression of disease symptoms of: Bean common mosaic virus (BCMV) Bean golden mosaic virus (BGMV) Beet curly top hybrigeminivirus (BCTV)	8.0 – 12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray at planting directed on or below seed; OR
3. In a narrow (2" or less) surface band over seed-line during planting incorporated to a depth of 1 to 1.5" with sufficient irrigation within 24 hours following application; OR
4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; OR
5. As a post-seeding drench, transplant-water drench, or hill drench.

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

FOLIAR APPLICATIONS (FIELD)

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	1.4

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum amount of product allowed per crop season: **4.2 fluid ounce** (0.13 lb AI/A)

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

ROOT VEGETABLES: Beet (garden)[†], Burdock (edible)[†], Carrot[†], Celeriac[†], Chervil (turnip-rooted)[†], Chickory[†], Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip[†], Radish[†], Oriental radish (diakon)[†], Rutabaga[†], Salsify (black)[†], Salsify (oyster plant), Salsify (Spanish), Skirret, Turnip[†]

TUBEROUS and CORM VEGETABLES (except Potato): Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet)[†], Chayote (root), Chufa, Dasheen (taro)[†], Ginger, Leren, Sweetpotato, Tanier (cocoyam)[†], Turmeric, Yam bean (jicama, manioc pea), Yam (true)[†]

SOIL APPLICATIONS¹ (FIELD)

Pests	Fluid ounces/1,000 row ft.	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Thrips (foliage-feeding) Whiteflies	0.35 – 0.85	5.0 – 12.0

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. In-furrow spray (rate specified per 1,000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting; OR
3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting; OR
4. Side-dress not more than 0.6 fluid ounces/1000 row-feet no later than 45 days after planting.

Remarks

Important Note: The rate applied affects the length of control. Use higher specified rates within the rate range where infestations occur later in crop development, or where pest pressure is continuous. Rates of this product less than 0.7 fluid ounce/1,000 row-feet will not provide adequate residual pest control. Crops treated with this product grown on very high organic matter soils (muck) may also require additional pest management control.

[†] The tops or greens from these crops may be utilized for food or feed.

Restrictions

Pre-Harvest Interval (PHI) for Root Vegetables: **21 days**

Pre-Harvest Interval (PHI) for Tuberous and Corm Vegetables: **3 days** (leaves); **125 days** (corms)

Maximum amount of product allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Maximum applications per crop season: **1**

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

¹ Soil application into Root Vegetables is not permitted in California unless otherwise directed by state approved 24(c) labeling.

(continued)

ROOT VEGETABLES: *(continued)***TUBEROUS and CORM VEGETABLES (except Potato):** *(continued)***FOLIAR APPLICATIONS¹ (FIELD)**

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	1.4
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Remarks	
† The tops or greens from these crops may be utilized for food or feed.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days <u>Maximum amount of product allowed per crop season:</u> Radish: 1.4 fluid ounce (0.044 lb AI/A) All other listed crops: 4.2 fluid ounces/Acre (0.13 lb. AI/A) <u>Maximum applications per crop season:</u> Radish: 1 All other listed crops: 3 DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling. ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	

SUGARBEET: (California only)**SOIL APPLICATIONS (FIELD)**

Pests	Fluid ounces/Acre
For control of: Aphids Flea beetles Leafhoppers Whiteflies	3.0 – 6.0
For suppression of disease symptoms of: Western yellows virus Beet curly top hybrigeminivirus (BCTV)	

(continued)

SUGARBEET: (California only) *(continued)***Application Methods**

Apply specified dosage of this product in the following method:

Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting.

Remarks

The low rate may be applied to aid establishment of stands in whitefly areas, or for early season control of the other pests listed.

Restrictions

Maximum amount of product allowed per year: **6.0 fluid ounces/Acre** (0.19 lb AI/Acre)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

DO NOT use on crops grown for seed unless permitted by state approved 24(c) labeling.

NUPRID® 4F INSECTICIDE CONVERSION CHART FOR LINEAR APPLICATION ONLY										
RATE fluid ounces/ Acre	RATE fluid ounces/1,000 row-feet Based on <u>average</u> row spacing (in inches):									
	10	15	20	25	30	34	36	38	40	45
5.0	0.10	0.14	0.19	0.24	0.29	0.33	0.34	0.36	0.38	0.43
5.5	0.11	0.16	0.21	0.26	0.32	0.36	0.38	0.40	0.42	0.47
6.0	0.11	0.17	0.23	0.29	0.34	0.39	0.41	0.44	0.46	0.52
6.5	0.12	0.19	0.25	0.31	0.37	0.42	0.45	0.47	0.50	0.56
7.0	0.13	0.20	0.27	0.33	0.40	0.46	0.48	0.51	0.54	0.60
7.5	0.14	0.22	0.29	0.36	0.43	0.49	0.52	0.55	0.57	0.65
8.0	0.15	0.23	0.31	0.38	0.46	0.52	0.55	0.58	0.61	0.69
8.5	0.16	0.24	0.33	0.41	0.49	0.55	0.59	0.62	0.65	0.73
9.0	0.17	0.26	0.34	0.43	0.52	0.59	0.62	0.65	0.69	0.77
9.5	0.18	0.27	0.36	0.45	0.55	0.62	0.65	0.69	0.73	0.82
10.0	0.19	0.29	0.38	0.48	0.57	0.65	0.69	0.73	0.77	0.86
10.5	0.20	0.30	0.40	0.50	0.60	0.68	0.72	0.76	0.80	0.90
11.0	0.21	0.32	0.42	0.53	0.63	0.72	0.76	0.80	0.84	0.95
11.5	0.22	0.33	0.44	0.55	0.66	0.75	0.79	0.84	0.88	0.99
12.0	0.23	0.34	0.46	0.57	0.69	0.78	0.83	0.87	0.92	1.03
12.5	0.24	0.36	0.48	0.60	0.72	0.81	0.86	0.91	0.96	1.08
13.0	0.25	0.37	0.50	0.62	0.75	0.85	0.90	0.95	0.99	1.12
13.5	0.26	0.39	0.52	0.65	0.77	0.88	0.93	0.98	1.03	1.16
14.0	0.27	0.40	0.54	0.67	0.80	0.91	0.96	1.02	1.07	1.21

Important Note: Rate of this product applied affects the length of control and, to a considerable extent, the degree of control or effect. Row-spacing X rate combinations in shaded blocks may not provide adequate residual pest control and are not recommended for long-term, residual control. Use higher labeled rates where infestations may occur later in crop development or where pest pressure is continuous. Nufarm offers no warranty for use of this product at rates below 0.35 fluid ounce/1,000 row- feet (the Row-Spacing/Rate combinations that are shaded).

BERRY, BUSH and VINE CROPS

STRAWBERRY: annual and perennial varieties	
SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Whiteflies	12.0 – 16.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening; OR 2. As a plant material or plant hole treatment just prior to, or during transplanting; OR 3. As a pre-plant band spray over the row in a minimum of 20 gallons of water per acre, followed immediately by overhead irrigation to incorporate product into root zone. DO NOT use plastic or other mulch that limits movement of this product into root zone.	
Remarks	
The rate applied affects the length of control. Use higher specified rates within the rate range where infestations may occur later in crop development or where pest pressure is continuous.	
Restrictions	
Pre-Harvest Interval (PHI): 14 days Maximum amount allowed per crop season: 16.0 fluid ounces/Acre (0.50 lb AI/Acre) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. DO NOT make both a soil and foliar application on the same crop in the same season. DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.	
SOIL APPLICATIONS (post-harvest use on perennial varieties)	
Pests	Fluid ounces/Acre
For control of: White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	8.0 – 12.0
Application Methods	
Apply a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply specified dosage of this product in one of the following methods: 1. As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre; OR 2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth must be equivalent to the width of the anticipated fruiting bed; OR 3. As a chemigation application with 600 to 1,000 gallons of water followed by 0.10 to 0.25 inch irrigation.	

STRAWBERRY: annual and perennial varieties <i>(continued)</i>	
Remarks	
All soil-surface applications must be followed by 0.25 inch of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate this product into egg-deposition zone may result in decreased activity of beetle grubs.	
Restrictions	
Maximum amount allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A)	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Spittlebugs Whiteflies	1.5
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Remarks	
All soil-surface applications must be followed by 0.25 inch of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate this product into egg-deposition zone may result in decreased activity of beetle grubs.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between applications: 5 days Maximum amount of product allowed per crop season: 4.6 fluid ounces/Acre (0.14 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. DO NOT make both a soil and foliar application on the same crop in the same season. DO NOT use on crops grown for seed unless allowed by state approved 24(c) labeling.	

BUSHBERRY: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal	
SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Japanese beetle (adults, feeding on foliage) White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. 18-inch band on each side of the row followed with 0.25 inch of irrigation immediately after application.	
Remarks	
For grub control, apply this product to control 1st or 2nd (early) instar larvae. Application may be made post-bloom up to 7 days prior to harvest, or post-harvest until October 1st. For control of Japanese beetle larvae, make applications from June 1 to July 15. DO NOT apply during bloom. Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root zone will help protect berry plant roots from grub feeding. Apply this product to moist soil. If necessary, apply one hour of irrigation water immediately before application. To ensure maximum efficacy, 0.5 to 1 inch of irrigation water or rainfall must be applied or received within 24 hours of application of this product to facilitate movement into the soil and into the root zone.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb AI/Acre) DO NOT apply pre-bloom or during bloom or when bees are foraging.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters	1.2 – 1.6
For control of: Japanese beetles (adults) Thrips (foliage feeding)	2.4 – 3.2
For control of: Blueberry maggot	3.2

continued

BUSHBERRY: (continued)**Application Methods**

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **3 days**

Minimum interval between applications: **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

Maximum number of product applications per year: **5**

Maximum application volume (water): **Ground: 20.0 GPA; Air: 5.0 GPA**

DO NOT apply pre-bloom or during bloom or when bees are foraging.

CANEBERRY: Blackberry (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these). Raspberry (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*).

SOIL APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	8.0 – 16.0
For control of: Rednecked cane borer	12.0 – 16.0
For suppression of: Thrips (foliage-feeding only)	16.0

Application Methods

Apply specified dosage in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. Basal, soil drench in a minimum of 500 gallons solution per acre.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

(continued)

CANEBERRY: *(continued)*

FOLIAR APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers	3.2
For suppression of: Thrips (foliage-feeding only)	
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 3 days Minimum interval between applications: 7 days Maximum amount of product allowed per season: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are actively foraging.	
¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	

CRANBERRY

SOIL APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Rootgrubs (Scarab) Rootworms (Chrysomelid)	8.0 – 16.0
Application Methods	
Apply this product to moist soil. Apply specified dosage of this product in one of the following methods: 1. As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre; OR 2. As a chemigation application with 600 to 1,000 gallons water. Immediately upon application, this product must be incorporated into root zone by 0.1 to 0.3 inch water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control. Make application post-bloom immediately after honeybees are removed. Application should target early instar larvae.	

(continued)

CRANBERRY *(continued)***Remarks**

Best control may be achieved when application is made post-bloom immediately after bees are removed. Target early instar larvae. This product has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of this product and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.

Restrictions

Pre-Harvest Interval (PHI): **30 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

GRAPES: American bunch grape, Muscadine grape, and Vinifera grape.

SOIL APPLICATIONS

Pests	Fluid ounces/Acre
For control of: European fruit lecanium Leafhoppers/Sharpshooters Mealybugs Phylloxera spp. ¹	8.0 – 16.0
For suppression of: Grapeleaf skeletonizer Nematodes ²	16.0
For suppression of disease symptoms of : Pierce's disease	

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation; OR
3. Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation.

(continued)

GRAPES: (continued)**Remarks**

Make application between bud-break and the pea-berry stage. A total of 14 fluid ounces/acre is required under the following conditions:

1. Where vigorous vine growth is expected
2. In warmer growing areas
3. Where mealybug and European fruit lecanium populations are expected to be heavy
4. Where vine populations exceed 600 per acre, or;
5. For suppression of nematodes

¹ Repeated and regular use of this product over multiple consecutive growing seasons controls existing *Phylloxera* infestations over time or prevents *Phylloxera* from becoming established.

² For suppression of nematodes, apply 14 fluid ounces in a single application or two 7-fluid ounce applications on a 30 to 45-day interval. Only make treatments by 1) chemigation into root zone through above ground low pressure drip, tickle, micro sprinkler or equivalent equipment or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root zone of the plant. Repeated and regular use of this product over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

Restrictions

Pre-Harvest Interval (PHI): **30 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Leafhoppers/Sharpshooters Mealybugs	1.2 – 1.6
For control of: Grapeleaf skeletonizer	1.6

Application Methods

Apply specific dosage of this product using properly calibrated ground application equipment only. Apply as a broadcast or directed spray to infested areas ensuring thorough coverage.

Restrictions

Pre-Harvest Interval (PHI): **0 days**

Minimum interval between applications: **14 days**

Maximum amount of product allowed per year: **3.2 fluid ounces/Acre** (0.1 lb. AI/A)

HOPS:

SOIL APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids	9.6
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR 2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation; OR 3. Hill drench in sufficient water to insure incorporation into the root zone followed by irrigation.	
Restrictions	
Pre-Harvest Interval (PHI): 60 days Maximum amount of product allowed per year: 9.6 fluid ounces/Acre (0.3 lb AI/Acre) ¹ This use is not permitted in CA unless otherwise specified by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids	3.2
Application Methods	
Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.	
Restrictions	
Pre-Harvest Interval (PHI): 28 days Minimum interval between applications: 21 days Maximum amount of product allowed year: 9.6 fluid ounces/Acre (0.3 lb. AI/A)	

COFFEE:

SOIL APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Leafminers	8.0 – 16.0
For suppression of: Scales	

(continued)

COFFEE: (continued)**Application Methods**

Apply specified dosage in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.; OR
2. Subsurface side-dress shanked into the root zone on both sides of the plants followed by irrigation.; OR
3. Basal, soil drench in sufficient water to insure incorporation into the root zone followed by irrigation.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Whiteflies	3.2
For suppression of: Scales	

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

CITRUS, TREE NUT and ORCHARD CROPS

CITRUS (containerized): Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

SOIL APPLICATIONS

Pests	mL/ft³ of container
For control of: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies	0.38
For control of: Citrus root weevil (larval complex) ¹	0.63 – 1.25
For suppression of : Citrus thrips (foliage-feeding only)	1.25

Application Methods

Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of this product per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, treatment must be made at planting prior to insect infestation. Retreat if necessary.

Application Restrictions

Pre-Harvest Interval (PHI): **0 day**
Maximum allowed per application: **0.50 mLs/0.1 ft³ container media**
Maximum allowed per crop season: **3.0 mLs/plant**
Do not apply pre-bloom or during bloom or when bees are foraging.

Remarks

¹ For control of larvae of the citrus root weevil complex, make application prior to neonate larvae entering potting media. Utilize specified higher dosage for heavy infestations.

CITRUS: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

SOIL APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Termites (FL only) Whiteflies	8.0 – 16.0
For suppression of: Citrus nematode Thrips (foliage-feeding thrips only)	16
For suppression of disease symptoms of: Citrus tristeza virus (CTV) through vector control Citrus yellows	

Application Methods

Apply specified dosage of this product in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. Apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil must be lightly pre-wetted to break soil surface tension prior to applications of this product. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move this product into root zone. Allow 24 hours before initiating subsequent irrigations; OR
2. Soil surface band spray on both sides of the tree. Bands must overlap at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root zone. This method is suitable for very coarse soils with 0.75% organic matter or less; OR
3. Drench to base of tree not exceeding one quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only for trees up to 8 feet tall; OR
4. For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk; OR
5. For suppression of citrus nematode, apply specific dosage through low pressure chemigation or soil surface spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of this product over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

(continued)

CITRUS: *(continued)***Restrictions**Pre-Harvest Interval (PHI): **0 days**Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)**DO NOT** apply during bloom or within 10 days prior to bloom or when bees are foraging.**FOLIAR APPLICATIONS**

Pests	Fluid ounces/100 gallons	Fluid ounces/Acre
For control of: Aphids Asian citrus psyllid Black fly Leafhoppers/Sharpshooters Leafminers Mealybugs Scales ¹ Whiteflies	1.4 – 2.0 (dilute application)	4.0 – 8.0 (dependent on tree size, target pest, and infestation pressure)
For suppression of: Thrips (foliage-feeding thrips only)	2.0	8.0

Application Methods

Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply through properly calibrated ground or aerial equipment.

Remarks

Aerial application of this product may result in slower activity and reduced control compared to ground application. Where higher rate applications are appropriate, increase the spray solution concentration to apply an equivalent rate per acre to that applied in the diluted application. The 8.0 fluid ounce/Acre rate is based on full sized trees. This rate may be reduced proportionally for smaller trees.

¹ Scales - time applications to the crawler stage. Treat each generation.

RestrictionsPre-Harvest Interval (PHI): **0 days**Minimum interval between sprays: **10 days**Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)**DO NOT** apply during bloom or within 10 days prior to bloom or when bees are foraging.

POME FRUIT: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince		
SOIL APPLICATIONS		
Pests	Fluid ounces/Acre	
For control of: Aphids (including Woolly apple aphid) Leafhoppers	8.0 – 12.0	
Application Methods		
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.		
Restrictions		
Pre-Harvest Interval (PHI): 21 days Maximum amount of product allowed per year: 12.0 fluid ounces/Acre (0.38 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.		
FOLIAR APPLICATIONS		
Pests	Fluid ounces/100 gallons	Fluid ounces/Acre
For control of: Leafhoppers	0.4 – 0.8	1.6 – 3.2
For control of: Aphids (except Woolly apple aphid) Apple maggot Leafminers San Jose scale	0.8	3.2
For use on Pears Only to control: Mealybugs Pear psylla	2.0	8.0
Application Methods		
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.		
Remarks		
Combine applications targeting apple maggot with manufacturer's specified rate of a sticker.		
Restrictions		
Pre-Harvest Interval (PHI): 7 days Minimum interval between sprays: 10 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.		

POMEGRANATE	
SOIL APPLICATIONS ¹	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Whiteflies	8.0 – 16.0
Application Methods	
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 0 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging. ¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	
FOLIAR APPLICATIONS	
Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Whiteflies	3.2
For suppression of: Scales	
Application Methods	
Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 7 days Minimum interval between sprays: 7 days Maximum amount of product allowed per year: 9.6 fluid ounces/Acre (0.3 lb. AI/A) DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.	

STONE FRUIT: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

PRE-PLANT ROOT DIP APPLICATIONS

Pests	Fluid ounces/10 gallons root dip solution
For control of: Black peach aphid (infesting roots)	1.0

Application Methods

Mix this product at a rate of 1.0 fluid ounce per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in this product's solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.

SOIL APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids (including Wooly apple aphid) Leafhoppers	8.0 – 12.0

Application Methods

Apply specified dosage of this product in the following method: Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum amount of product allowed per year: **12.0 fluid ounces/Acre** (0.38 lb. AI/A)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

FOLIAR APPLICATIONS

Pests	Fluid ounces/100 gallons	Fluid ounces/Acre
For control of: Aphids Green June beetle Japanese beetle Leafhoppers/Sharpshooters Plant bugs Rose chafer San Jose scale	0.8	1.6 – 3.2
For control of: Cherry fruit fly (maggot of Eastern & Western)		2.4 – 3.2
For suppression of: Plum curculio Stinkbugs		3.2

STONE FRUIT: *(continued)***Application Methods**

Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.

Restrictions**Apricot, Nectarine, Peach:**

Pre-Harvest Interval (PHI): **0 days**

Minimum interval between applications: **7 days**

Maximum amount of product allowed per crop season: **9.6 fluid ounces/Acre** (0.30 lb. AI/A)

Minimum application volume (water): **Ground: 50 GPA; Air: 25 GPA**

DO NOT apply during bloom or within 10 days prior to bloom or when bees are actively foraging.

Cherry, Plum, Plumcot, Prune:

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **10 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb. AI/A)

Minimum application volume (water): **Ground: 50 GPA; Air: 25 GPA**

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

TREE NUTS except Almonds: Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)**SOIL APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Mealybugs Spittlebugs Termites Two-lined spittlebugs Whiteflies	8.0 – 16.0
For suppression of: Thrips (foliage-feeding only)	16.0
For suppression of disease symptoms of: Pecan scab (from reduction in honeydew deposition)	

(continued)

TREE NUTS (except Almonds): *(continued)***Application Methods**

Apply specified dosage prior to or at onset of pest infestation in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. Pre-wet soil prior to applications of this product and allow soil to dry following application and prior to subsequent irrigation; OR
2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site; OR
3. Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Apply this product in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area must follow within 48 hours to promote uptake by root system; OR
4. For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18 to 24 inches. Allow soil to dry following treatment and prior to applying any irrigation.

Remarks

Use higher specified rates within the rate range when applied by shank or subsurface side-dress, used on larger trees, soils with high clay content, for high plant populations, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.

Restrictions

DO NOT apply in Almonds

Pre-Harvest Interval (PHI): **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb. AI/A)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

¹ Soil Application in Tree Nut orchards is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids (except Black pecan aphid) Leafhoppers/Sharpshooters Phylloxera spp. (leaf infestations) Spittlebugs Whiteflies	1.4 – 2.8
For control of: Black pecan aphid Mealybugs San Jose scale ¹	3.2

(continued)

TREE NUTS (except Almonds): *(continued)***Application Methods**

Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.

Remarks

¹ Applications for control of San Jose scale must be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.

Restrictions

DO NOT apply in Almonds

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **6 days**

Maximum amount of product allowed per year: **11.5 fluid ounces/Acre** (0.36 lb. AI/A)

Minimum application volume (water): **Ground: 50 GPA; Air: 25 GPA**

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

BANANA and PLANTAIN**SOIL APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers	8.0 – 16.0
For suppression of: Scales	
Application Methods	
Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.	
Restrictions	
Pre-Harvest Interval (PHI): 0 days	
Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb. AI/A)	
¹ This use is not permitted in CA unless otherwise directed by state approved 24(c) labeling.	

(continued)

BANANA and PLANTAIN *(continued)***FOLIAR APPLICATIONS**

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers Thrips	3.2
Application Methods	
Apply specified dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.	
Remarks	
Aerial application of this product may result in slower activity and reduced control relative to results from ground application. Addition of an organosilicone adjuvant at a rate not to exceed 2.0 fluid ounces/100 gallons finished spray solution may improve coverage and pest control.	
Restrictions	
Pre-Harvest Interval (PHI): 0 days Minimum interval between applications: 14 days Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.5 lb. AI/A)	

TROPICAL FRUIT: Acerola, Atemoya¹, Avocado, Biriba¹, Black sapote, Canistel, Cherimoya¹, Custard apple¹, Feijoa, Jaboticaba, Guava, Llama¹, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop¹, Spanish lime, Star apple, Starfruit, Sugar apple¹, Wax jambu.

SOIL APPLICATIONS¹

Pests	Fluid ounces/Acre
For control of: Aphids Avocado lacebug Leafhoppers Whiteflies	12.0 – 16.0
For suppression of: Scales Thrips (foliage-feeding thrips only)	16.0

(continued)

TROPICAL FRUIT: *(continued)***Application Methods**

Apply specified dosage of this product in the following method: Chemigation into root-zone through low-pressure (drip, trickle, micro-sprinkler or equivalent) equipment.

Restrictions

Pre-Harvest Interval (PHI): **6 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb. AI/A)

¹ Soil application use on noted crops is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Leafhoppers/Sharpshooters Mealybugs Thrips Whiteflies	3.2
For suppression of: Thrips	

Application Methods

Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.

Remarks

Ground applications of this product are more effective than aerial applications.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **10 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

DO NOT apply during bloom or within 10 days prior to bloom or when bees are foraging.

OTHER CROPS**Christmas Trees****SOIL APPLICATIONS¹**

Pests	Fluid ounces/Acre
For control of: White grub complex (e.g., grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0

Application Methods

Soil incorporation and movement of this product to the root zone is required for activity. This product can be incorporated most readily when applied to moist soil. Apply specified dosage in one of the following methods:

1. Chemigation into root zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; OR
2. 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25 to 1 inch of irrigation within 12 hours after application.

Remarks

Apply this product during adult flight activity, or up to mid-July, when first instar larvae are present.

Restrictions

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.50 lb. AI/A)

¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS

Pests	Fluid ounces/Acre
For control of: Aphids Adelgids Sawflies	1.6 – 3.2

Application Methods

Apply specific dosage of this product as a broadcast or directed spray to infested area ensuring thorough coverage. Apply this product through properly calibrated ground or aerial equipment.

Remarks

Ground applications of this product are more effective than aerial applications.

For gall-forming adelgids, time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. Once galls form spraying this product is ineffective.

Restrictions

Minimum interval between applications: **7 days**

Maximum amount of product allowed per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

Poplar/Cottonwood: (including members of the genus <i>Populus</i> grown for pulp or timber)	
SOIL APPLICATIONS¹	
Pests	Fluid ounces/Acre
For control of: Aphids Cottonwood leaf beetle	8.0 – 16.0
For suppression of: <i>Phylloxera popularia</i>	
Application Methods	
Apply specified dosage of this product in one of the following methods: 1. Chemigation through low-pressure drip irrigation; OR 2. For narrow row, cutting orchards/nurseries used for plant propagation, shank into root zone followed by adequate irrigation to promote uptake. Adequate irrigation depends on soil moisture level at application. Under dry conditions 0.25 inch/acre is recommended.	
Remarks	
For Cottonwood leaf beetle, protection against damage will occur when application is made early-season, when beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake. For <i>Phylloxera</i> , apply early in the year, from break of dormancy through May.	
Restrictions	
Maximum amount of product allowed per year: 16.0 fluid ounces/Acre (0.50 lb. AI/A) ¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.	
CUTTING/WHIP APPLICATIONS¹	
Pests	Soaking Solution Fluid ounces needed per 100 gallons
For control of: Cottonwood leaf beetle	6.65 to 13.3 (unhydrated cuttings/whips) 13.3 to 20.0 (partially hydrated cuttings/whips)
For suppression of: Aphids <i>Phylloxera popularia</i>	13.3 (unhydrated cuttings/whips) 20.0 (partially hydrated cuttings/whips)
Application Methods	
Apply this product in one of the following cuttings/whips soaking methods: 1. For freshly cut (hydrated) cuttings/whips, soak plant material in specified solution concentration for 24 hours prior to cold storage. After removal from cold storage, plant as needed; OR 2. For previously hydrated cuttings/whips removed from cold storage, allow plant material to reach room temperature and soak in specified solution concentration for 24 hours prior to planting. Take proper care in disposal of any residual soaking solution. Apply solution to existing trees or other registered crops as long as all product label precautions and restrictions are observed.	

Poplar/Cottonwood: (continued)**Remarks**

The moisture content prior to application of the cuttings/whips, the solution concentration and the length of soaking interval interact to affect the amount of product absorbed into plant material. For a constant soaking interval of 24 hours, dry cuttings/whips absorb a higher quantity of solution and require a lower concentration. Conversely, more hydrated cuttings/whips absorb less solution and require a higher concentration. Soaking of cuttings/whips must occur in a covered container in absence of UV light. Not all *Populus* sp. clones/varieties/hybrids have been tested for crop safety. Without specific knowledge about a particular *Populus* sp. clone/variety/hybrid, a small number of cuttings/whips of each must be treated and evaluated prior to commercial use.

Restrictions

Maximum amount of product allowed at plant per year: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

¹ This use is not permitted in California unless otherwise directed by state approved 24(c) labeling.

FOLIAR APPLICATIONS¹

Pests	Fluid ounces/Acre
For control of: Aphids Leaf beetles	1.6 – 3.2

Application Methods

Apply this product as a broadcast or directed spray method through properly calibrated ground, aerial or chemigation application equipment. Thorough coverage of foliage is necessary.

Remarks

Ground application of this product is more effective than aerial application for these crops.

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **10 days**

Maximum amount of product allowed per season: **16.0 fluid ounces/Acre** (0.5 lb. AI/A)

DO NOT apply during bloom or with 10 days prior to bloom or when honeybees are foraging.

¹ Use as a foliar application to Poplar/Cottonwood is not permitted in California unless otherwise directed by state approved 24(c) labeling.

COMMERCIAL POULTRY FACILITIES**POULTRY HOUSING STRUCTURES**

Pests	Fluid ounces / 1,000 ft²
For control of: Darkling beetles Hide beetles (Dermestids)	3.0 (90 ml)
Application Methods	
<p>Apply between flocks after de-caking and sanitation procedures have been completed. Apply as a spot, crack and crevice, or surface spray on floors, walls, and support beams of structure. Apply using a minimum of 1/2 to 2 gallons of spray mixture per 1000 square feet. To prepare the spray mixture, fill the spray tank with 1/2 the required amount of water, then add the specified amount of product. Add the remaining water while agitating or mixing. Maintain constant agitation while applying.</p> <p>Apply spray mixture to the entire footing including 1 foot up the wall above the footing, and in 3 to 4 foot wide bands directly beneath all feed lines. The areas beneath the feed lines typically harbor large numbers of adult and larval stages of the target pest when an infestation occurs. Measure these areas to determine the appropriate amount of spray mixture to apply. For structures that are prone to large infestations, treat the footings including 1 foot up the wall and the entire floor area of the structure.</p> <p>Cracks and crevice areas also are prone to large infestations of the target pest. Apply as a crack and crevice treatment around wall insulation or other areas that may harbor the target pest. If structures have supporting beams, treat the floor with a 1 foot band around each beam and apply 2 feet up the beam.</p> <p>For structures prone to extreme infestation, treat the entire structure with a broadcast application. Apply 3.0 fluid ounces in 2 gallons of water per 1000 square feet of surface. Apply as a broadcast spray to areas where litter has accumulated (floor, under feed and water lines, lower sections of walls, corners).</p>	
Remarks	
In order to avoid problems with pest resistance to imidacloprid, rotate to an insecticide with a different mode of action every 2-3 flocks. Rotate between 3 different insecticide mode of action classes labeled for control of target pests during a calendar year.	
Restrictions	
<p>DO NOT apply when birds are present or within 7 days of bird placement.</p> <p>DO NOT allow food or feed to be contacted by the spray.</p> <p>Remove feed and water from the treatment area before applying.</p>	
Pests	Fluid ounces / Gallon
For control of: Nuisance ants	0.125 – 0.25 (3/4 – 1.5 TSP)

(continued)

POULTRY HOUSING STRUCTURES *(continued)***Application Methods**

Apply as a crack and crevice or wall void treatment inside structures. Apply to cracks, crevices, drilled holes, onto walls, around entry points such as doors, windows, vents, eaves, soffits, and utility access openings. If nests are present in voids, apply into the void if possible. Apply evenly to treatment surfaces but not to the point of runoff. Apply to areas around the exterior of the structure where ants may be present (soil, turf, ornamental shrubs and plantings, and groundcover in close proximity to or touching the structure). For above-ground nests, such as in wood posts, decks, or fences, or in trees, spray into holes/openings where ants are traveling and on the wood surface.

Restrictions

DO NOT use for control of native or imported fire ants, harvester ants or pharaoh ants. Keep people and pets out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

NUPRID is a registered trademark of Nufarm Americas Inc.

All other trademarks that appear on this label which are not owned by Nufarm Americas Inc. or its subsidiaries are the property of their respective owners.


(RV121213)



NOTES



59






NOTES



60



NUPRID® 4F MAX

INSECTICIDE

GROUP 4A INSECTICIDE

ACTIVE INGREDIENT:

Imidacloprid, 1 -[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 40.4%

OTHER INGREDIENTS:..... 59.6%

TOTAL: 100.0%

Contains 4 pounds of imidacloprid per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ATTACHED BOOKLET FOR COMPLETE FIRST AID AND PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.

For Medical Emergencies Only, Call (877) 325-1840.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid breathing spray mist. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid contact with eyes or clothing. Wear protective eye wear. Wear long sleeved shirt and long pants, socks, shoes, and chemical resistant gloves made of waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

Refer to booklet for complete Personal Protective Equipment, Engineering Control Statements, User Safety Recommendations, Environmental Hazards and Directions For Use.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. **DO NOT** walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

EPA Reg. No. 228-528

EPA Est. No. indicated by the first two letters of the batch number on this package

(AL) 228-IL-002,

(YU) 67545-AZ-001

Net Contents: 1 Gal (3.78 L)

Manufactured for Nufarm Americas Inc.

11901 S. Austin Avenue | Alsip, IL 60803

PULL HERE TO OPEN

12815000

(RV121213)

file: 48209-4_base_art
folder: 48209-4_Nufarm_12815000
colors: black
☐ dp proofed

created: 12-13-07 jw
alt: 05-15-14 jw
size: 5.5"(w) x 5"(h)



Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803
Phone: 708-377-1330 Fax: 708-377-1333

4020 Aerial Center Parkway, Suite 101
Morrisville, NC 27560
Phone: 919-379-2510 Fax: 919-467-5923
www.us.nufarm.com

Via Courier Delivery

May 27, 2014

Ms. Venus Eagle, PM #1
USEPA (7504P)
Document Processing Desk (NOTIF)
Room S4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

**Subject: Application for NOTIFICATION(s) per PRN 98-10
Nuprid 2F, EPA Reg. No. 228-484;
Nuprid 4.6F Pro, EPA Reg. No. 228-527; and
Nuprid 4F Max, EPA Reg. No. 228-528**

Dear Ms. Eagle:

Enclosed please find three (3) application(s) for pesticide registration – NOTIFICATION for the above-referenced product(s). The purpose of these NOTIFICATION(s) is to correct the rate discrepancies for citrus (containerized). The rates were originally taken from *Admire Pro*, EPA Reg. No. 264-827, which is a 4.6 lb ai/gal product, but were not adjusted accordingly. These NOTIFICATION(s) are consistent with PRN 98-10. In support of these NOTIFICATION(s) enclosed are the following:


- EPA form 8570-1 Application for Pesticide Registration – NOTIFICATION; and
- Proposed product labeling in PDF format on CD-rom.

Should you have any questions or require additional information, please contact me at (919) 379-2528 or by email at carrie.tackema@us.nufarm.com.

Sincerely,


Carrie M. Tackema
Regulatory Manager

Enclosure(s)



Material Sent for Data Extraction

Reg. # 228-528

Description: CSF Amendment

☐ Material(s) Sent to Data Extraction Contractors:

☐ New Stamped Label Dated _____

☐ Notification Dated _____

☐ New CSF(s) Dated 10/7/14

☐ Other: _____

☒ Decision #: 497329

☐ Other Action/Comments: _____

File this coversheet and attached materials in the jacket. It must be well organized and clipped together, NOT STAPLED. Then give the jacket with the coversheet and materials to staff in the Information Services Center (ISC) (Room S-4900). If a jacket is full or only available as an image, please file materials in a new jacket and bring it down to the (ISC). For further information please call 703-605-0716.

Reviewer: Jessica Rosales

Phone: 347-0263 Division: RD/IRB

Date: 11/27/14